

Opinion

***Saccharomyces cerevisiae*: a nomadic yeast with no niche?**

5

Matthew R Goddard^{1,2} and Duncan Greig^{3,4}

¹*The School of Biological Sciences, The University of Auckland, Auckland 1142, New*

10

Zealand

²*The School of Life Sciences, The University of Lincoln, Lincoln LN6 7DL, United*

Kingdom

³*Max Planck Institute for Evolutionary Biology, Plön 24306, Germany*

⁴*Department of Genetics, Evolution, and Environment, University College London,*

15

London WC1E 6BT, United Kingdom

m.goddard@auckland.ac.nz; d.greig@ucl.ac.uk

20

Running head: *S. cerevisiae* neutral model

20

Abstract:

Different species are usually thought to have specific adaptations, which allow them to occupy different ecological niches. But recent neutral ecology theory suggests that species diversity can simply be the result of random sampling, due
25 to finite population sizes and limited dispersal. Neutral models predict that species are not necessarily adapted to specific niches, but are functionally equivalent across a range of habitats. Here we evaluate the ecology of *S. cerevisiae*, one of the most important microbial species in human history. The artificial collection, concentration, and fermentation of large volumes of fruit for
30 alcohol production produces an environment in which *S. cerevisiae* thrives, and therefore it is assumed that fruit is the ecological niche that *S. cerevisiae* inhabits and has adapted to. We find very little direct evidence that *S. cerevisiae* is adapted to fruit, or indeed to any other specific niche. We propose instead a neutral nomad model for *S. cerevisiae*, which we believe should be used as the
35 starting hypothesis in attempting to unravel the ecology of this important microbe.

Keywords:

Crabtree effect, fermentation, fruit, adaptation, niche, neutral ecology, natural
40 history

One sentence summary:

It is assumed that *S. cerevisiae* is adapted to inhabit fruits; however, we find very little evidence for adaptation to any niche. Instead we propose a neutral nomad

45 model for *S. cerevisiae*.

Adaptation and the ecological niche

The concept of a niche is central to the field of ecology. This concept presumes that there are specific sets of environmental conditions under which different species can thrive, and therefore that there are discrete places and times in
50 which species may be found. Classical ecological theory is in line with this concept and suggests that different species have different sets of “functions”, which they acquired under adaptation by natural selection to different ecological niches (Vandermeer 1972). Fundamental constraints restrict the number of functions a single species can have – an organism cannot simultaneously enjoy
55 the benefits of being both big and small for example. Such trade-offs explain biodiversity: any given habitat supports multiple species because no single species can successfully occupy all niches. The metaphor that species have specific functions that allow them to occupy specific niches was developed by direct observation of macroscopic species, and on the whole it satisfactorily
60 explains both biodiversity and why species appear to fit their environments (Vandermeer 1972). This idea has been transferred to the microbial world, where the possibility of huge population sizes and high rates of dispersal should increase the power of natural selection to drive adaptation to specific niches (Hanson *et al* 2012). This thinking is epitomised by the Baas Becking hypothesis
65 (Baas Becking 1934): “everything is everywhere, but the environment selects”. However, direct observations of microbial interactions with their natural environments are often impossible, and it may be that the ecological niche concept is not generally applicable to microbes.

Neutral theory of ecology

Recently, other models have been proposed that successfully explain species diversity. Neutral ecology emphasizes the importance of stochastic processes in determining community structure and function (Bell 2001; Hubbell 2005). These
75 models present diversity as the result of random sampling, caused by finite population sizes and limited dispersal. This idea implies that species may not be preferentially adapted to different niches, and that in fact different species might be functionally equivalent across a number of niches. Neutral models of ecology remain controversial and have not yet been widely applied to microbial ecology
80 (Hanson *et al* 2012), except for examples in which limited dispersal is seen to be a primary determinant of community composition (e.g. Peay *et al.* 2010; Bell 2010; Dumbrell *et al* 2010). Here we reconsider whether the ecological niche metaphor applies to the yeast *Saccharomyces cerevisiae*, one of the best studied laboratory model organisms, but whose ecology and natural history is still
85 largely unknown. We present the idea that *S. cerevisiae* is not adapted to a specific niche, but is instead a nomad that has evolved the general ability to inhabit and persist in many different environments.

The importance of *Saccharomyces cerevisiae*

90 *S. cerevisiae* has been widely used by humans for thousands of years and is arguably one of the most important microbial species in human history (Chambers *et al.* 2010). It owes this distinction to a single trait: its ability to produce alcohol from sugar. Whilst it is also useful for raising bread, producing fuel, and expressing desirable engineered proteins, it was the demand for
95 alcoholic beverages that motivated the scientific study of yeast by Pasteur

(1897) and the Carlsberg Research Laboratories (Hansen 1896). Since then *S. cerevisiae* has achieved a second distinction: it is the best understood genetic model organism. *S. cerevisiae* was the first eukaryote to have its genome completely sequenced, and its genome is still the best annotated and most tractable to genetic manipulations and analysis (Cherry 2011). Huge projects are in the process of determining the biological functions and genetic interactions of every part of the genome (e.g. Kelly *et al.* 2001; Boone 2014) on a scale that is unprecedented in any organism. *S. cerevisiae* has been key to numerous major breakthroughs in genetics, biochemistry, and cell biology (Chambers *et al.* 2010).

105

The Crabtree Effect

S. cerevisiae preferentially produces alcohol from sugar by anaerobic fermentation, even when oxygen is available for aerobic respiration. This key trait, known as the Crabtree effect (Pronk *et al.* 1996), is thought to be an adaptation to high sugar environments. Although fermentation of sugar by *S. cerevisiae* is about ten times less metabolically efficient than aerobic respiration in terms of ATP production, it potentially provides two proposed selective benefits. First, fermentation liberates energy faster and thus enables more rapid growth than aerobic respiration does (Pfeiffer *et al.* 2001). If many individuals compete for a limited shared resource, those that grow more rapidly will win, even if they effectively squander the resource (Pfeiffer *et al.* 2001; MacLean *et al.* 2006). A useful metaphor for this is “the tragedy of the commons” (Hardin 1968). Secondly, fermentation degrades the environment by producing ethanol, which is not produced by aerobic respiration. In addition fermentation produces heat and CO₂ more rapidly than aerobic respiration does, so these may also

120

accumulate. If *S. cerevisiae* can tolerate such alcoholic, hot, and anoxic environments better than its competitors, then it will enjoy a selective advantage due to the interference effects of its own fermentation, and there is some experimental evidence to support this idea (Goddard 2008). Although often seen
125 as competing hypotheses, these two potential benefits of the Crabtree effect are not mutually exclusive but complementary. Further, having outgrown or interfered with its competitors, *S. cerevisiae* can then undergo a 'diauxic shift' and switch metabolic gears to use the accumulated ethanol as a substrate for aerobic respiration, recovering some (but not all) of the energy wasted by
130 fermentation (Thomson *et al.* 2005). This reduction of the metabolic cost of initial fermentation is available as a consequence of either or both of the two earlier benefits, but it is usually associated with the second, in the so-called "make-accumulate-consume" strategy (Piskur *et al.* 2006).

135 The Crabtree effect is thought to have originated around the time that the ancestor of the *Saccharomyces* clade underwent a whole genome duplication (Piskur *et al.* 2006). Whilst most duplicated gene copies were subsequently lost, many of the surviving genes play roles in sugar metabolism and may have been maintained because two copies allow increased relative gene expression (Kellis
140 *et al.* 2004). Further, the presence of two copies of a gene allows one to maintain ancestral function, whilst the other is free to diverge and acquire new functions. One such case appears to be the duplicated gene pair *ADH1* and *ADH2* (Thomson *et al.* 2005). *ADH1* reduces acetaldehyde to ethanol during anaerobic respiration. *ADH2*, though, appears to have diverged so that it catalyzes the reverse reaction
145 underpinning the diauxic shift: it reconverts ethanol to acetaldehyde, which can

be used to make Acetyl-CoA, which feeds into the citric acid cycle. This neofunctionalisation has been proposed as one of the key innovations underlying the Crabtree effect, allowing *S. cerevisiae* both to tolerate ethanol and to recover energy that would otherwise be wasted by fermentation. The fact that the
150 Crabtree effect is thought to have appeared at around the same time as fruiting plants became widespread is cited as evidence that *S. cerevisiae* is adapted to a specific niche, fruit. However, the evidence is circumstantial: dating the origin of the Crabtree effect in geological time using only genetic data is very error prone, and more recent work suggests that the Crabtree effect may have evolved over a
155 long period of time, and is not just coincidental with the whole genome duplication (Hagman *et al.* 2013).

Does the superiority of *S. cerevisiae* as a fermenter indicate that it is adapted to fruit?

160 The best evidence that yeast are adapted to fruit comes from winemaking. When grapes are gathered and crushed, they spontaneously ferment, producing wine (e.g. Goddard 2008). Given the prominence of alcohol in human history (McGovern *et al.* 2004), this basic process must have worked fairly consistently and reliably for a long time, across a wide range of conditions. As *S. cerevisiae* is
165 the primary microbe associated with winemaking, it appears logical to assume that its natural habitat includes grapes and the other fruits that are used to make alcoholic beverages.

However, the natural fruit habitat differs greatly from the artificial fermentation
170 environment created by makers of wine and other traditional alcoholic

beverages. Collecting large numbers of individual fruits and crushing them together homogenises the resource and increases its size, making it equally accessible to all species present. By mixing many individual communities from many individual fruits together into one, the number and diversity of individuals competing increases, and this creates competitive conditions that may favour high rate, rather than high efficiency, of growth (MacLean *et al.* 2006). Physical containment within a vessel and the decrease in surface area to volume ratio of the sugar resource might enhance the degradation of the environment by preventing ethanol, CO₂, and heat from dissipating easily as it might from individual fruits in the open, selecting for the interference effects of fermentation. It will also reduce the ability of oxygen to diffuse into the fruit, selecting for fermentation. Winemaking conditions are therefore expected to favour fermentation much more strongly than the conditions on natural fruit, so the success of *S. cerevisiae* in wine does not therefore imply that it is successful on fruit.

The adaptation model predicts that organisms adapted to a niche should be abundant in that niche. But *S. cerevisiae* is in fact vanishingly rare on fruit, even in vineyards where fruiting plants are artificially at very high densities and the associated winemaking would be expected to increase the overall abundance of yeast in the location (Mortimer *et al.* 1999; Knight *et al.* 2014). Metagenomic sequencing suggests *Saccharomyces sp.* comprises less than 1:20,000 of the fungi on ripe grapes in vineyards; instead, Crabtree-negative yeast species dominate (Taylor *et al.* 2014). Other yeast species initially dominate early fermentation of wine, and only after several days of fermentation does *S. cerevisiae* typically

become abundant (Goddard 2008; Ciani *et al.* 2010). Indeed, other species may often persist at significant frequency even in the extremely alcoholic conditions at the end of wine fermentation (Goddard 2008; Jolly *et al.* 2014). Further, despite artificial conditions that strongly favour fermentation, it is common for
200 spontaneous wine ferments to get 'stuck' – that is no fermentative microbe dominates and very little ethanol is produced (Bisson *et al.* 2000).

The artificial nature of the conditions that are used to ferment wine and other alcoholic drinks is emphasised by the need for humans to produce alcohol
205 themselves, rather than to collect it from some natural source. Whilst anecdotal reports of animals getting drunk on naturally occurring alcohol are common, well-known examples such as elephants (Morris *et al.* 2006) and waxwings (Eriksson *et al.* 1983) have been debunked. In addition, although low levels of ethanol can occur in fruits (Eriksson *et al.* 1983) and nectar (Wiens *et al.* 2008),
210 there is no evidence that *Saccharomyces* is the primary microorganism responsible. Thus, although *S. cerevisiae* tends to become the dominant organism when large numbers of fruit are gathered, combined, and fermented by winemakers, it does not therefore follow that *S. cerevisiae* is well adapted to fruits under natural conditions. Indeed, given its scarcity on fruits, even in
215 vineyards, it seems reasonable to question whether *S. cerevisiae* is especially adapted to fruit at all.

Is the Crabtree effect a spandrel?

In evolutionary biology, a spandrel is a trait that exists as the by-product of the
220 evolution of some other trait, rather than because it was the target of selection.

The word refers to architectural spandrels, triangular areas of masonry between structural arches supporting a dome. These alluringly-shaped spaces are often highly decorated, and it is tempting to view them as the main feature around which the rest of the building is designed. But this is not so: spandrels exist
225 merely as a necessary by-product of a dome supported by arches. In a classic paper Gould and Lewontin (1979) used spandrels as an analogy to persuade evolutionary biologists to not view all organisms' traits as the product of adaptation by natural selection. The Crabtree effect might appear to us to be an important adaptation because of the reverence with which we regard the
230 fermentation of wine, but it is possible that the Crabtree effect did not evolve as an adaptation to ferment natural fruit, or is even as an adaptation at all.

The proposed benefits of the Crabtree effect – rapid growth and interference effects on competitors – have not been quantified experimentally in the natural
235 fruit environment, only in homogenised grape juice in the laboratory (Goddard 2008), or in artificial media (MacLean *et al.* 2006). Even in fermenting wine, the anti-competitor benefit of ethanol production by *S. cerevisiae* is modest (~2%), and some yeast species found in spontaneous ferments are not suppressed by ethanol until it exceeds 9%, i.e. close to the end of fermentation (Goddard 2008).
240 Further, fermentation by *S. cerevisiae* occurs even in sugar concentrations 100-fold less than those typically found in fruit (Pfieffer *et al.* 2014). Such low glucose levels would be expected to select for high growth efficiency rather than high growth rate, and the interference effects of fermentation would be negligible at these sugar levels as miniscule levels of ethanol, heat and CO₂ would be
245 produced. It seems quite possible, therefore, that fermentation has benefits other

than high growth rate and interference. For example, perhaps the ethanol serves as an attractant for insect vectors, rather than as an anti-competitor compound (Buser *et al.* 2014). The idea that the Crabtree effect is an adaptation enabling competitive superiority under conditions resembling the artificial winery environment might therefore be a spandrel. Even if the Crabtree effect can be shown to be advantageous under some condition, we cannot easily assess whether natural selection shaped it unless we can assess how often yeast encounters similar conditions in nature.

It is also possible that the Crabtree effect is not a fixed trait in *S. cerevisiae*; however, that the Crabtree effect appears invariant across closely related species make this possibility unlikely. The reason this is a possibility at least is that all *S. cerevisiae* (and for that matter all *Saccharomyces sensu stricto*) have been isolated due to their ability to ferment and gain competitive superiority in sugar-rich environments. This occurred either during spontaneous wine ferments or via enrichment culture isolation procedures, which essentially mimic the winemaking process: typically, environmental samples such as pieces of plant material are placed in a sugary liquid medium and incubated at an elevated temperature. Enrichment cultures are often spiked with ethanol, to favour ethanol-resistant genotypes. Thus, the isolation of *S. cerevisiae* from natural samples itself selects for Crabtree positive strains, and would likely leave Crabtree negative strains undetected. Variation in fermentation ability and alcohol tolerance clearly exists even among the current biased sample of *S. cerevisiae* isolates (Stern 2014), and is therefore likely to be greater in nature.

270 Thus, there may well be Crabtree negative, or at least less strongly Crabtree
positive, *Saccharomyces* genotypes in nature of which we are unaware.

What is *S. cerevisiae*'s niche?

As we have stated: the adaptation model predicts that organisms should be
275 abundant in niches to which they are adapted. The belief that *S. cerevisiae* is
adapted to inhabit fruits derives from its dominance in the spontaneous
fermentations of wine. But, as we argue above, the considerable difference
between the winery environment and natural fruits, combined with the low
abundance of *S. cerevisiae* on fruits, and especially on wine grapes, suggests that
280 this belief may be incorrect. Another feature of fruit is that it is an ephemeral
resource. What then does *S. cerevisiae* do when it has exhausted a fruit of
nutrients? The simplest explanation is that it disperses to another fruit, perhaps
passively or perhaps by an insect vector. Recent work shows that some *S.*
cerevisiae release volatile compounds that attract *Drosophilid* flies, which can
285 vector it from fruit to fruit (Buser *et al.* 2014). But what happens when the
fruiting season is over? Diploid yeast cells may undergo meiosis and turn into
more resistant haploid spores when starved, so it is possible that once a fruit
resource is exhausted, large numbers of yeast cells sporulate and disperse as
dormant, resistant spores which might persist for many years, allowing them to
290 survive not only the dispersal process, but also the long intervals of time
between fruiting seasons. Consistent with this model are the observations that
yeast spores, but not vegetatively growing cells, are resistant to *Drosophila*
digestion (Reuter *et al.* 2007), and that *S. cerevisiae* has been isolated from

Drosophila (Buser *et al.* 2014), hibernating wasps (Stefanini *et al.* 2014), and bee
295 hives (Goddard *et al.* 2010).

This raises the possibility that *S. cerevisiae* is adapted to habitats other than fruit,
and these may form a refuge when fruit is not available. Finding and
characterising these putative niches is an active field of research. *S. cerevisiae*
300 has been isolated from a wide range of environments: in addition to vineyard
and winery environments and a range of other human ferments (sake, billi wine,
etc.) and baking (Liti *et al.* 2009), *S. cerevisiae* is found, as expected, in fruits and
insects, but also in humans as a commensal (Angebault *et al.* 2013) or pathogen
(Muller *et al.* 2011), in soil, on various plants (Wang *et al.* 2012), and on oak
305 trees (Sniegowski *et al.* 2002; Sampaio *et al.* 2008). The belief that oak is another
of *S. cerevisiae*'s niches comes from a seemingly consistent ability to isolate yeast
from oak bark. Unfortunately it is likely that this survey is biased: many
researchers simply want samples of wild yeast to study, and therefore they
target environments from which *Saccharomyces* has already been isolated. Since
310 oak trees are typically sympatric with yeast laboratories, and are easy for
microbiologists to identify, it is not surprising that they are a favoured source of
wild yeast samples.

The enrichment culture method that is nearly always used to isolate yeast also
315 likely causes severe biases, not only in favour of Crabtree positive yeast, as
discussed above, but also in obscuring the true distribution and range of *S.*
cerevisiae. If a sample does not yield *S. cerevisiae* by enrichment culture, it does
not mean that it was not present or viable, but only that it did not grow

sufficiently to outcompete the other microbes in the sample. Nor is it possible to
320 tell whether any *S. cerevisiae* strains that are recovered originated from dormant
spores in the sample, or from an actively growing population. The observed
homozygosity of soil and wild oak-tree associated *Saccharomyces* might
therefore be an artefact of the isolation process, if isolates are derived from rare
single spores that auto-diploidise in the enrichment culture by mating-type
325 switching (Goddard *et al.* 2010). Consistent with this explanation, mitotic
diploids isolated directly from wine ferments are typically much more
heterozygous than those from oak bark and soil (Goddard *et al.* 2010; Knight *et*
al. 2014).

330 Simple enrichment culture does not give any indication of the abundance of *S.*
cerevisiae in a primary sample. By determining the sensitivity of enrichment
culture to detect single yeast cells spiked into oak bark samples, and making
appropriate dilutions of oak bark samples, we have estimated the average
density of *Saccharomyces paradoxus*, *S. cerevisiae*'s sister species, to be just two
335 cells per square centimetre of oak bark; consistent with this extremely low
density, we detected no *Saccharomyces* sequences at all among 40,000 fungal
sequences extracted from oak bark (Kowallik *et al.*, *in revision*). We are unaware
of such estimates for *S. cerevisiae*. Further, recent work shows that some *S.*
cerevisiae isolates grow poorly in 'oak bark extract' in the laboratory (Giraldo-
340 Perez *et al.* 2013). Together, the evidence suggests that oak bark is not a niche
that *S. cerevisiae* is especially abundant in or well-adapted to.

Genetic variation within *S. cerevisiae*

The global population structure of *S. cerevisiae* shows evidence for clades
345 associated with specific environments, consistent with adaptation of certain
genotypes to certain environments (Liti *et al.* 2009). But there is also an equal
weight of evidence for isolation by distance, consistent with neutral divergence
due to limited dispersal and subsequent lack of gene flow between
geographically remote populations (Liti *et al.* 2009; Knight *et al.* 2014). Limited
350 sampling of such wild populations by humans making artificial fermentations
could result in the association between genotype and environment appearing by
chance, rather than by adaptation. The prime case is for the population
inhabiting Europe. Humans inadvertently amplified this sub-population with
their winemaking, and provided an increased density of opportunities for *S.*
355 *cerevisiae* at some point ~9,000 years ago when we began to deliberately grow
fruit and make wine in one place (Le Gras *et al.* 2007). This Wine/European
lineage is now no longer geographically constrained because humans have
transported it around the globe along with viticulture and wine making (Le Gras
et al. 2007; Liti *et al.* 2009; Goddard 2010; Gayevskiy & Goddard *in prep*). Whilst
360 this *S. cerevisiae* lineage now has one apparent adaptation to agricultural
interventions, resistance to copper and sulphur that are used as anti-microbials
in vineyards (Aa *et al.* 2006), there is no other compelling evidence that the
founders of this lineage were differentially adapted or better at fermenting than
individuals from other lineages. A possible way to determine whether wild
365 populations are adapted to specific habitats is to use a form of reciprocal
transplant experiment. For example, a set of strains isolated from different
habitats could be tested in direct competition assays in conditions simulating the
different habitats. If strains tend to have high relative fitness in the environment

they were isolated from but low relative fitness elsewhere, it would indicate that
370 they were indeed adapted to specific conditions.

The Nomad model

The ripe speculation in these last paragraphs indicates our lack of data
concerning *S. cerevisiae's* natural history. We have little direct evidence
375 concerning the niche or niches that *S. cerevisiae* might be abundant in or adapted
to. We know little about its distribution, or the form it takes in different habitats
(dormant spores or vegetative cells); nor do we know how the asexual, sexual,
and dormant phases of its life cycle fit into its life history. *S. cerevisiae* is known
to be abundant only in the ferments of artificially gathered fruit, but it appears to
380 be sparsely distributed everywhere else that has been surveyed for, and
particularly sparse on fruits and oak tree bark, the most commonly claimed
niches for this species.

This leads us to propose an alternative neutral model: that *S. cerevisiae* is not
385 adapted to a specific niche, but is a nomad, able to survive as a generalist at low
abundance in a wide range of environments. This is consistent with the
observation that *S. cerevisiae* can be found in a diverse range of habitats. The low
abundance of this species is also consistent with this it being a generalist,
capable of doing lots of things, but none of them especially well. *S. cerevisiae* has
390 a rich metabolism that enables it to survive or grow in a wide range of
environments eclipsing those found in either fruits or bark, with varying:
nutrient availabilities - with both low and high carbon and nitrogen
concentrations (Gray *et al.* 2012; Wenger *et al.* 2011); pH - from strongly acidic

~pH 3 (Goddard 2008) to alkaline ~pH8 (Serrano *et al.* 2006); osmolarity -
395 from survival in water through to NaCl concentrations of at least 1.3M
(Petrovska *et al.* 1999); and temperatures ranging from close to zero to around
45 °C (Salvado *et al.* 2011; Sweeny *et al.* 2004). These observations suggest the
fundamental niche, the set of conditions where *S. cerevisiae* may survive, is very
broad, and this is in line with it being a generalist. Comprehensive fitness
400 measurements across a range of conditions (including interactions with other
species), for a wide range of genotypes, may indicate the realised niche for *S.*
cerevisiae: the range of conditions actually used by this species. Such a task is not
trivial however. Lastly, there is genomic evidence that *S. cerevisiae* is a
generalist, not a specialist: the genome is complex, containing 6,000 genes, of
405 which only 20% are necessary for growth in simple laboratory medium (Giaever
et al. 2002).

S. cerevisiae's diverse metabolic tolerances, range of habitats of isolation, and low
densities are therefore consistent with a nomad model, but they are not
410 inconsistent with it being adapted to some other, as yet undetermined niche or
niches. The main utility of the Nomad Model, then, is as a neutral scientific
starting point, which can serve as a null hypothesis for evaluating adaptive
explanations for *S. cerevisiae* evolution. Rather than assume that *S. cerevisiae* is
adapted to a niche, and then doggedly search for data that back up this prejudice,
415 we should start from the position of neutrality.

Future methods that could be used to test proposed niche models against the
neutral Nomad Model include experimental evolution in a candidate niche to test

whether genes are maintained by selection in a given environment such as grape
420 juice, or whether superfluous pathways are lost, indicating that the tested
environment is not a niche that *S. cerevisiae* is adapted to. Unbiased surveys –
that sample niches systematically, regardless of where *S. cerevisiae* has
previously been discovered – will provide data as to the incidence of this species
in various habitats. The increasing torrent of environmental sequence data will
425 also provide opportunities for realistic estimates for *S. cerevisiae*'s abundances.
The Nomad Model can be rejected if comprehensive, systematic, and unbiased
surveys reveal that *S. cerevisiae* inhabits and is competitively robust in specific
natural habitats and not others. Predictable changes in abundance indicating
seasonal growth cycles would also be valuable evidence supporting adaptation.
430 Such data could explain how the life cycle of *S. cerevisiae* fits in with the changes
in its natural environment.

Conclusion

435 It is tempting to apply the niche adaptation concept to *S. cerevisiae*, and to
microbes in general, as this concept is familiar to us by observation of large
organisms. However, we must be cautious not to assume adaptation to specific
niches, but demand evidence for such adaptations. The nomad model serves as a
neutral model for *S. cerevisiae*, and is the counterpoint to the assertion that the
440 species is adapted to one or more specific niches. We believe it should be used as
the null hypothesis for research attempting to unravel the ecology of this
important model microbe.

445 *Acknowledgements*

This idea was jointly conceived at the ISSY31 meeting in Vipava, Slovenia, and then discussed, developed, and written up by the authors when they were Visiting Fellows of the Isaac Newton Institute for Mathematical Sciences, Cambridge, England.

450

Literature cited

- Aa E, Townsend JP, Adams RI, Nielsen KM, Taylor JW. Population structure and gene evolution in *Saccharomyces cerevisiae*. *FEMS Yeast Res* 2006; 6:702–715.
- 455 Angebault C, Djossou F, Abélanet S, *et al.* *Candida albicans* Is Not Always the Preferential Yeast Colonizing Humans: A Study in Wayampi Amerindians. *Journal of Infectious Diseases* 2013; 208: 1705–1716.
- Baas Becking L. *Geobiologie of inleiding tot de milieukunde*. W.P. Van Stockum & Zoon, The Hague, 1934.
- 460 Bell G. Neutral macroecology. *Science* 2001; 293, 2413-2418.
- Bell T. Experimental tests of the bacterial distance–decay relationship. *The ISME Journal* 2010; 4: 1357–1365.
- Bisson LF and Butzke CE. Diagnosis and Rectification of Stuck and Sluggish Fermentations. *Am. J. Enol. Vitic.* 2000; 51: 168-177.
- 465 Boone C. Yeast Systems Biology: Our Best Shot at Modeling a Cell. *Genetics* 2014; 198: 435–437.
- Buser CC, Newcomb RD, Gaskett AC, Goddard MR. Niche construction initiates the evolution of mutualistic interactions. *Ecology Letters* 2014; 17(10):1257-1264.
- 470 Chambers PJ, Pretorius IS. Fermenting knowledge: the history of winemaking, science and yeast research. *EMBO Rep.* 2010; 11: 914–920.
- Cherry JM, Hong EL, Amundsen C, *et al.* *Saccharomyces* Genome Database: the genomics resource of budding yeast. *Nucleic Acids Research* 2011; 40:D700–D705.
- 475 Ciani M, Comitini F, Mannazzu I, Domizio P. Controlled mixed culture fermentation: a new perspective on the use of non-*Saccharomyces* yeasts in

- winemaking. *FEMS Yeast Res.* 2010; 10(2):123-33.
- 480 Dumbrell AJ, Nelson M., Helgason T *et al.* Relative roles of niche and neutral processes in structuring a soil microbial community. *The ISME Journal* 2010; 4, 337-345.
- Eriksson K and Nummi H. Alcohol accumulation from ingested berries and alcohol metabolism in passerine birds. *Ornis Fennica* 1983; 60: 2–9.
- Gayevskiy V, Goddard MR. Phylogenomic signals quantify patterns of human-aided *S. cerevisiae* trans-hemisphere range expansion. *In prep.*
- 485 Giaever G, Chu AM, Ni L, *et al.* Functional profiling of the *Saccharomyces cerevisiae* genome. *Nature* 2002; 418, 387–391.
- Giraldo-Perez P, Goddard MR. A parasitic selfish gene that affects host promiscuity. *Proc. R. Soc. B* 2013; 280 20131875.
- 490 Goddard MR. Quantifying the complexities of *Saccharomyces cerevisiae*'s ecosystem engineering via fermentation. *Ecology* 2008; 89: 2077–2082.
- Gould SJ, Lewontin, R C. The spandrels of San Marco and the Panglossian paradigm: a critique of the adaptationist programme. *Proc. R. Soc. London B: Biol. Sci.* 1979; 205, 581-598
- 495 Hagman A, Sall T, Compagno C, Piskur J. Yeast “Make-Accumulate-Consume” Life Strategy Evolved as a Multi-Step Process That Predates the Whole Genome Duplication. *PLoS ONE* 2013; 8.
- Hanson CA, Fuhrman JA, Horner-Devine MC, Martiny JBH. Beyond biogeographic patterns: Processes shaping the microbial landscape. *Nature Reviews Microbiology* 2012; 10, 497-506
- 500 Hansen E C. *Practical studies in fermentation being contributions to the life history of micro-organisms.* E & FN Spon, London, 1896.
- Hardin G. Tragedy of Commons. *Science* 1968; 162: 1243–&.
- Hubbell SP. Neutral theory in community ecology and the hypothesis of functional equivalence. *Functional Ecology* 2005; 19, 166-172.
- 505 Jolly NP, Varela C, Pretorius I S. Not your ordinary yeast: non-*Saccharomyces* yeasts in wine production uncovered. *FEMS Yeast Research* 2014;. 14(2): 215–237.
- Kelly DE, Lamb DC, Kelly SL. Genome-wide generation of yeast gene deletion strains. *Comparative and Functional Genomics* 2001; 2: 236–242.
- 510 Kowallik, Miller, Greig D. The interaction of *Saccharomyces paradoxus* with its natural competitors on oak bark. *Molecular Ecology*; accepted pending minor revisions.

- 515 Knight SJ, Goddard MR. Quantifying separation and similarity in a *Saccharomyces cerevisiae* metapopulation. *The ISME Journal* 2014; Advance online publication 25 July 2014.
- Legras J-L, Merdinoglu D, Cornuet J-M, Karst F. Bread, beer and wine: *Saccharomyces cerevisiae* diversity reflects human history. *Molecular Ecology* 2007;16:2091–2102.
- 520 Liti G, Carter DM, Moses AM, et al. Population genomics of domestic and wild yeasts. *Nature* 2009; 458:337–341.
- MacLean RC, Gudelj I. Resource competition and social conflict in experimental populations of yeast. *Nature* 2006; 441: 498-501.
- McGovern PE, Zhang J, Tang J *et al.* Fermented beverages of pre-and proto-historic China. *PNAS* 2004; 101:17593–17598.
- 525 Morris S, Humphreys D, Reynolds D. Myth, Marula, and Elephant: An Assessment of Voluntary Ethanol Intoxication of the African Elephant (*Loxodonta africana*) Following Feeding on the Fruit of the Marula Tree (*Sclerocarya birrea*). *Physiol Biochem Zool* 2006;79: 363–369.
- 530 Mortimer R, Polsinelli M. On the origins of wine yeast. *Res Microbiol* 1999; 150: 199–204.
- Muller LAH, Lucas JE, Georgianna DR, McCusker JH. Genome-wide association analysis of clinical vs. nonclinical origin provides insights into *Saccharomyces cerevisiae* pathogenesis. *Molecular Ecology* 2011; 20: 4085–4097.
- 535 Otterstedt K, Larsson C, Bill R *et al.* Switching the mode of metabolism in the yeast *Saccharomyces cerevisiae*. *EMBO Rep* 2004; 5: 532–537.
- Pasteur L. *Studies on fermentation: the diseases of beer, their causes, and the means of preventing them.* Macmillan & Co., London, 1897.
- 540 Peay KG, Garbelotto M, Bruns TD. Evidence of dispersal limitation in soil microorganisms: Isolation reduces species richness on mycorrhizal tree islands. *Ecology* 2010; 91: 3631–3640.
- Petrovska B, Winkelhausen E, Kuzmanova S. Glycerol production by yeasts under osmotic and sulfite stress. *Can. J. Microbiol* 1999; 45: 695–699.
- 545 Pfeiffer T, Schuster S, Bonhoeffer S. Cooperation and competition in the evolution of ATP-producing pathways. *Science* 2001; 292, 504–507.
- Pfeiffer T, Morley A. An evolutionary perspective on the Crabtree effect. *Front. Mol. Biosci.* 2014; 1:17
- Piskur J, Rozpedowska E, Polakova S, Merico A, Compagno C. How did *Saccharomyces* evolve to become a good brewer? *Trends Genet* 2006; 22:

- 550 183–186.
- Pronk JT, Steensma HY, Van Dijken JP. Pyruvate metabolism in *Saccharomyces cerevisiae*. *Yeast* 1996; 12:1607-1633.
- Reuter M, Bell G, Greig D. Increased outbreeding in yeast in response to dispersal by an insect vector. *Current Biology* 2007; 17: R81–R83.
- 555 Sampaio JP, Gonçalves P. Natural Populations of *Saccharomyces kudriavzevii* in Portugal Are Associated with Oak Bark and Are Sympatric with *S. cerevisiae* and *S. paradoxus*. *Applied and Environmental Microbiology* 2008; 74(7):2144–2152
- 560 Serrano R, Martín H, Casamayor A and Ariño J. Signaling Alkaline pH Stress in the Yeast *Saccharomyces cerevisiae* through the Wsc1 Cell Surface Sensor and the Sit2 MAPK Pathway. *The Journal of Biological Chemistry* 2006; 281, 39785-39795
- 565 Salvadó Z, Arroyo-López FN, Guillamón JM *et al.* Temperature Adaptation Markedly Determines Evolution within the Genus *Saccharomyces*. *Appl. Environ. Microbiol.* 2011; 77(7): 2292-2302
- Sniegowski PD, Dombrowski PG, Fingerman E. *Saccharomyces cerevisiae* and *Saccharomyces paradoxus* coexist in a natural woodland site in North America and display different levels of reproductive isolation from European conspecifics. *Fems Yeast Res* 2002; 1(4):299-306.
- 570 Stern DL. Identification of loci that cause phenotypic variation in diverse species with the reciprocal hemizyosity test. *Trends in Genetics* 2014; 30 (12):547–554.
- Stefanini I, Dapporto L, Legras, JL *et al.* Role of social wasps in *Saccharomyces cerevisiae* ecology and evolution. *PNAS* 2012; 109: 13398–13403.
- 575 Sweeney JY, Kuehne HA, Sniegowski PD. Sympatric natural *Saccharomyces cerevisiae* and *S. paradoxus* populations have different thermal growth profiles. *FEMS Yeast Research* 2004; 4: 521-525
- 580 Taylor MW, Tsai P, Anfang N, Ross HA, *et al.* Pyrosequencing reveals regional differences in fruit-associated fungal communities. *Environmetnal Microbiology* 2014; 16(9): 2848–2858.
- Thomson JM, Gaucher EA, Burgan MF, *et al.* Resurrecting ancestral alcohol dehydrogenases from yeast. *Nat Genet* 2005; 37: 630–635.
- Vandermeer JH. Niche Theory. *Annual Review of Ecology and Systematics* 1972; 3: 107-132.
- 585 Wang Q-M, Liu W-Q, Liti G, *et al.* Surprisingly diverged populations of *Saccharomyces cerevisiae* in natural environments remote from human activity. *Molecular Ecology* 2012; 21(22):5404-17.

Wenger JW, Piotrowski J, Nagarajan S, *et al.* Hunger artists: yeast adapted to carbon limitation show trade-offs under carbon sufficiency. *PLoS Genet.* 2011; 7(8):e1002202.

Wiens F, Zitzmann A, Lachance M *et al.* Chronic intake of fermented floral nectar by wild treeshrews. *s* 2008; 105(30):10426-31.