

A short proof of the Gröttsche conjecture.

Joint with M Kool + V Shende.

- Steiner 1848 } \mathbb{P}^2
- Cayley } \mathbb{P}^2
- Salmon } \mathbb{P}^2
- \vdots
- Ran, Choi \mathbb{P}^2
- Vainsencher S
- Di Francesco - Itzykson \mathbb{P}^2
- Caporaso-Harris \mathbb{P}^2
- You-Zaslow (Beauville, Bryan-Lung) $K3$
- Gröttsche S
- Kleiman-Piene S
- A. Liu (S, w)
- Kazarian S
- Fomin - Mikhalkin \mathbb{P}^2
- Y. Tzeng 2010 S


$p(x,y)$ degree n poly in 2 complex variables

Zero locus is complex curve

$$\{p(x,y)=0\} \subseteq \mathbb{C}^2$$

Generically smooth

modelled on copy of \mathbb{C} $\{\underline{v} : \underline{v} \cdot \nabla p = 0\}$ near zeros of p where $\nabla p \neq 0$.



(i.e. local model just $x=0$)

Singular in codimension-1

3 eqns $p=0=\partial_x p=\partial_y p$
in 2 unknowns x,y


\Rightarrow In a generic 1-parameter family of curves,
linear, for simplicity

$$C_\lambda := \{p_\lambda := p_0 + \lambda p_\infty = 0\}$$

expect a finite number of simplest
singularities - nodes local model $\{xy=0\}$

$$+ \subseteq \mathbb{C}^2$$

(or $\{x^2+y^2=0\}$; p and first derivatives vanish,
 2^{nd} derivs nondegenerate)



Calculation with resultant of polys P_0, P_∞

$\Rightarrow (\leq) \underline{3(n-1)^2}$ of them.

(Will give 2 proofs shortly.) First, consider easier case; $P(x)=0$ in \mathbb{C} .

Generically smooth (n points).

Linear 1d family $P_\lambda(x) := P_0(x) + \lambda P_\infty(x) = 0$

Singular when get double root, when discriminant of P_λ vanishes.

↖ degree $2(n-1)$ in coefficients of P_λ
 \Rightarrow degree $2(n-1)$ in λ
 $\Rightarrow 2(n-1)$ such double points in our 1d linear family.


Proof via Euler characteristics:

Any $x \in \mathbb{P}^1$ is root of exactly one $P_\lambda = P_0 + \lambda P_\infty$

$(\mathbb{C} \cup \{\infty\})$

(assuming P_0, P_∞ have no common roots)

Gives map $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ n -fold cover.
 $x \mapsto \lambda$



Inverse image of $\lambda \in \mathbb{P}^1$ is $\begin{cases} n \text{ points} \\ (n-1) \text{ points} \end{cases}$

generically
at double
point of P_λ

Taking Euler characteristics gives

$$2 = n \cdot 2 - \#(\text{double points})$$


□

Back to $P_\lambda(x,y)$ - i.e. curves in \mathbb{P}^2 .

$$\deg P_\lambda = n \Rightarrow e(\{P_\lambda = 0\}) = 2 - (n-1)(n-2)$$

when $\{P_\lambda = 0\}$ smooth.

and $3 - (n-1)(n-2)$ when 1-nodal



each punctured smooth curve
has $e = 2 - (n-1)(n-2) - n^2$

$$e(\mathbb{P}^2) = e(\mathbb{P}^1) (2 - (n-1)(n-2) - n^2) + n^2 + \#(\text{nodal curves})$$

$$\Rightarrow \# \text{ nodal curves} = 3n^2 - 6n + 3 = 3(n-1)^2 \quad \square$$

General case.

S smooth complex surface

L sufficiently ample

↑ Gröttsche: $\geq (5\delta-1)$ -ample
Us: can get away with $\geq \delta$ -ample

$P(H^0(L)) \leftrightarrow$ curves $C \subset S$ st. $\mathcal{O}(C) = L$

- Generically smooth
- Singular in codim 1

generic
• 1-d family $\mathbb{P}^1 \subset P(H^0(L))$ has finite # of curves with only simplest possible singularities: nodes $\{xy=0\} \subset \mathbb{C}^2$
+

generic
• \hookrightarrow 2-d family $\mathbb{P}^2 \subset \mathbb{P}$ has finite #
of curves with 2 nodes (plus finite
number with cusps; 1-d family with 1 node)

generic
• \hookrightarrow δ -dim family $\mathbb{P}^\delta \subset \mathbb{P}$ has finite #
of δ -nodal curves (plus much else
besides; but crucially can prove
all other singular curves have
geometric genus $\bar{g} > g - \delta$).

How many?

Conjecturally topological, given by deg δ

Universal polynomial in $c_1(L)^2, c_1(L) \cdot c_1(S),$
 $c_1(S)^2, c_2(S).$

So deg 2δ poly in $n = \deg C$ for $(\mathbb{P}^2, \mathcal{O}(n))$

Eg 1-nodal curves in pencil $\mathbb{P}^1 \subseteq \mathbb{P}(H^0(L))$.

Approach #1

Section s of $L \otimes \mathcal{O}(1)$ on $S \times \mathbb{P}^1$.

Nodes \leftrightarrow simple zeros of (s, ds)

Section of $L \otimes \mathcal{O}(1) \oplus T_{S \times \mathbb{P}^1}^* \otimes (L \otimes \mathcal{O}(1))$

(or use 1-jets if you don't like connections)

\Rightarrow # nodal curves =

$$c_3(L \otimes \mathcal{O}(1) \oplus T_{S \times \mathbb{P}^1}^* \otimes (L \otimes \mathcal{O}(1)))$$


Calculate as $c_2(s) + 3c_1(L)^2 + 2c_1(s) \cdot c_1(L)$

Approach #2: Euler characteristics

Smooth curve C has

$$\begin{aligned} e &= 2 - 2g = -\deg_C(K_C) \\ &= -\deg_C(K_S \otimes L) \quad \text{by adjunction} \\ &= (c_1(S) - c_1(L)) \cdot c_1(L) \end{aligned}$$

Nodal curve has $e = (2 - 2g) + 1$.




S

$C \cdot C = c_1(L)^2$
points of intersection

each punctured smooth curve
has $e = 2 - 2g - c_1(L)^2$

$$e(S) = e(\mathbb{A}^1) (2 - 2g - c_1(L)^2) + c_1(L)^2 + \# \begin{matrix} \text{(nodal)} \\ \text{curves} \end{matrix}$$

Equivalently blow up S at $c_1(L)^2$ points



to get same formula

$$e(\hat{S}) = e(S) + c_1(L)^2 = e(\mathbb{P}^1)(2-2g) + \# \text{ (nodal curves)}$$

this is the universal curve \mathcal{C}
 \downarrow
 \mathbb{P}^1

$$e(\text{Hilb}^1(\mathcal{C}/\mathbb{P}^1)) = (2g-2)e(\text{Hilb}^0(\mathcal{C}/\mathbb{P}^1)) + \# \text{ (nodal curves)}$$

The key is the existence of an invariant of curves

$$e(C) - (2 - 2g) = \begin{cases} 0 & \text{if } C \text{ smooth} \\ 1 & \text{if } C \text{ nodal} \end{cases}$$

$= e(\text{Hilb}^1 C) - (2 - 2g)e(\text{Hilb}^0 C)$

"Integrating" (w.r.t. Euler characteristic) over the base gives #1-nodal curves.

More generally, for δ -nodal curves we have the following result
(joint work with R Pandharipande)

Thm [PT] \exists universal linear combination

$$n_{g-\delta}(c) := \sum_{i=0}^{\delta} \alpha_i e(\text{Hilb}^i C)$$

Such that, for all reduced curves of arithmetic genus g ,

$$n_{g-\delta}(c) = \begin{cases} 0 & \text{if geometric genus of } C \text{ is } > g-\delta \\ 1 & \text{if } C \text{ is } \delta\text{-nodal} \end{cases}$$

(GV BPS invt of C)

Inductively, $n_g(c) := 1$,

$$n_{g-1}(c) := e(c) - n_g(c)(2-2g),$$

$$n_{g-i}(c) := e(\text{Hilb}^i C) - \sum_{j=g-i+1}^g n_j(c) e(\text{Sym}^{i-(g-j)} \Sigma_j)$$

where $\Sigma_j =$ smooth curve of genus j

Intuitively, $n_i(C)$ is the number of curves of arithmetic genus i "carried by" C (eg. with a map to C which is generically an iso.)

Eg for δ -nodal C , get a genus $g-i$ curve mapping to it by resolving i nodes. $\binom{\delta}{i}$ such choices, and indeed

$$n_{g-i}(C) = \binom{\delta}{i} \text{ in this case}$$

"Integrating" (w.r.t. Euler characteristic)
over the base gives $\# \delta$ -nodal curves
as the same linear combination

$$\sum_{i=0}^{\delta} \alpha_i e(\text{Hilb}^i \mathcal{C}/\mathbb{P}^{\delta}).$$

So need to compute e of

$$\text{Hilb}^i \mathcal{C}/\mathbb{P}^{\delta}.$$

$$\text{Hilb}^i \mathbb{C}/\mathbb{P}^{\delta} \subseteq \text{Hilb}^i \mathbb{S} \times \mathbb{P}^{\delta}$$

Cut out by tautological section
of a tautological bundle E over
 $\text{Hilb}^i \mathbb{S} \times \mathbb{P}^{\delta}$.

$$\left\{ \begin{array}{l} \text{Fibre } E|_{(z, [s])} = H^0(L|_z) \otimes \mathcal{O}(1) \\ \text{Section has value } s|_z \end{array} \right.$$

$$\text{Vanishes} \Leftrightarrow s|_z = 0 \Leftrightarrow z \in C \quad \begin{array}{l} \text{curve } s^{-1}(0) \\ \swarrow \end{array}$$

Transverse! Zeros $\text{Hilb}^i \mathbb{C}/\mathbb{P}^{\delta}$ smooth of
codimension = $\text{rk } E$.

So we can write $e(\text{Hilb}^i \mathbb{P}^g)$ in terms of an integral of Chern classes of E and $T_{\text{Hilb}^i S \times \mathbb{P}^g}$ over $\text{Hilb}^i S \times \mathbb{P}^g$; and thus over $\text{Hilb}^i S$.

Ellingsrud - Göttsche - Lehn have an algorithm to turn such integrals into integrals over S^i .

\Rightarrow deg i poly in $c_1(L)^2, c_1(L) \cdot c_1(S), c_1(S)^2$ and $c_2(S)$.


\Rightarrow $n_{g-\delta}$ is deg δ poly in the same.

$\frac{c_2(S)^\delta}{\delta!}$ leading term

Computer code calculates only up to 4 nodes so far...
(correct answers though)

Motivation.

Obvious way to single out S -nodal curves is via maps from curves of genus $g-\delta$.




Count these via ("reduced") Gromov-Witten theory and GV BPS states. Hard!

MNOP conjecture
(really for 3-folds)

GW
GV \leftrightarrow "stable
pairs"

Count embedded curves with
free points on them ("D-branes")

Change in genus $g \mapsto g - \delta$ \leftrightarrow up to δ
free points



$g \mapsto g+1$
free pts $\uparrow +1$

To understand $e(\text{Hilb}^i)$ s of curves of arithmetic genus g , geometric genus $> g - \delta$, may as well remove smooth genus $g - \delta + 1$ bit and assume it's a rational curve of arithmetic genus $g = \delta - 1$.

Then AJ: $\text{Hilb}^i C \rightarrow \overline{\text{Pic}}_i C$
 has fibre $\mathbb{P}(H^0(L))$ over $L \in \overline{\text{Pic}}_i C$.
 $e = h^0(L)$ ($= 1 - g + i$ once $i > 2g - 2$)

$i = 0, 1, 2, \dots, g-1, g, g+1, \dots, 2g-2, 2g-1, \dots$

\nwarrow Serre dual \nearrow
 $L \mapsto L^* \otimes K_C$

e predetermined
 $(h^0(L) = 1 - g + i \text{ once } i > 2g - 2)$

All $e(\text{Hilb}^i C)$ s determined by those for $i \leq g$.

\Rightarrow Can write $e(\text{Hilb}^{g+1} C)$ as linear comb. of lower $e(\text{Hilb}^i C)$ s
 $e(\text{Hilb}^0) + \alpha_{\delta-1} e(\text{Hilb}^{\delta-1}) + \alpha_{\delta-2} \dots \equiv 0 \quad \forall C$ of geom. genus $> g - \delta$.
 Can check LHS applied to δ -nodal curve gives $+1$.