

MODULAR INVARIANT OF QUANTUM TORI

C. CASTAÑO BERNARD & T. M. GENDRON

ABSTRACT. The quantum modular invariant $j^{\text{qt}}(\theta)$ of $\theta \in \mathbb{R}$ is defined as a discontinuous $\text{PGL}_2(\mathbb{Z})$ -invariant multi-valued map using the distance-to-the-nearest-integer function $\|\cdot\|$. For $\theta \in \mathbb{Q}$ it is shown that $j^{\text{qt}}(\theta) = \infty$ and for quadratic irrationalities PARI/GP experiments suggest that $j^{\text{qt}}(\theta)$ is a finite set. In the case of the golden mean φ , we produce explicit formulas involving weighted versions of the Rogers-Ramanujan functions for the experimental supremum and infimum. We then define a universal modular invariant ${}^\circ j: {}^\circ\widehat{\text{Mod}} \rightarrow {}^\circ\hat{\mathbb{C}}$ as a *continuous* and *single valued* map of ultrasolenoids, such that 1) the classical modular invariant is a quotient of the restriction of ${}^\circ j$ to a subsolenoid $\text{Mod}^{\text{cl}} \subset {}^\circ\widehat{\text{Mod}}$ fibering over the classical moduli space of elliptic curves and 2) the quantum modular invariant is a quotient of the restriction of ${}^\circ j$ to a subsolenoid $\text{Mod}^{\text{qt}} \subset {}^\circ\widehat{\text{Mod}}$ fibering over the moduli space of elliptic curves equipped with a Kronecker foliation.

CONTENTS

Introduction	1
1. The Quantum Modular Invariant of a Real Number	4
2. Modular Invariant of the Golden Mean I: An Explicit Formula	7
3. Modular Invariant of the Golden Mean II: Convergence	12
4. Quantum Tori and Kronecker Foliations	15
5. Nonstandard Structures	17
6. Diophantine Approximation Groups	20
7. Ultrasolenoids	23
8. Eisenstein Ultratransversals	26
9. The Universal Modular Invariant	28
10. Appendix: Experimental Values of J^{qt}	32
References	35

INTRODUCTION

One of the most powerful links between number theory, geometry and analysis can be found in the theory of complex multiplication (CM): a signal event in a long development in number theory, which begins with Gauß's reciprocity law and culminates in the main theorems of class field theory [34], [36].

The theory of CM yields results that come under the heading of «explicit class field theory», which may be seen as generalizations of the Theorem of Kronecker-Weber. If $\omega \in \mathbb{C} - \mathbb{Q}$ is a complex quadratic irrationality, let $E_\omega(\mathbb{C}) \cong \mathbb{C}/\langle 1, \omega \rangle$ be the

Date: 31 January 2012.

2000 Mathematics Subject Classification. 11U10, 11F03, 14H52, 57R30, 11M55.

complex points of the elliptic curve E_ω parametrized by ω . Then $E_\omega(\mathbb{C})$ has CM (its endomorphism ring is strictly larger than \mathbb{Z}); if we denote $K = \mathbb{Q}(\omega)$ and assume that $\text{End}(E_\omega(\mathbb{C})) \otimes \mathbb{Q} = K$ then

- The maximal unramified abelian extension (Hilbert class field) H of K is generated over K by the modular invariant $j(\omega)$ of $E_\omega(\mathbb{C})$.
- The maximal abelian extension K^{ab} of K is generated over H by the values of the Weierstraß \wp -function at the torsion subgroup of $\mathbb{C}/\langle 1, \omega \rangle$.

Finding the analogue of the theory of CM for more general fields – Kronecker’s *Jugendtraum* or Hilbert’s twelfth problem – has been the focus of investigation for nearly a century [18], [35], [37], [23], [7].

In 2004, Yu. Manin [25] proposed an approach to the Stark conjectures [37] in which quantum tori play a role analogous to that of elliptic curves with CM. Manin’s *Alterstraum* is sometimes known as the «Real Multiplication» programme (RM): an approach to the Stark conjectures in the case of real quadratic fields which uses notions of noncommutative geometry.

In [25], the quantum torus $\mathbb{T}(\theta)$, $\theta \in \mathbb{R} - \mathbb{Q}$, is understood as an object in a category of irrational rotation C^* -algebras, however it may also be described in somewhat more naive geometric terms as the quotient $\mathbb{R}/\langle 1, \theta \rangle$ of the reals by the pseudo lattice $\langle 1, \theta \rangle$, or equivalently, as the space of leaves of the Kronecker foliation $\mathcal{F}(\theta)$ c.f. [27] or §4 of this paper. The moduli space of quantum tori is identified with the quotient $\text{Mod}^{\text{qt}} = \text{PGL}_2(\mathbb{Z}) \backslash (\mathbb{R} - \mathbb{Q})$.

One would therefore like to formulate and prove exact analogues of the main theorems of CM in the RM case, using suitable notions of Weierstraß \wp -function and modular invariant for the quantum torus $\mathbb{T}(\theta)$. However both $\mathbb{T}(\theta)$ and Mod^{qt} are quotients by groups acting with dense orbits, so in particular, there are no non constant continuous functions defined on either of them. Thus it is not at all clear how to define the analogues of the Weierstraß \wp -function or modular invariant in this setting.

The goal of the present paper is to provide a definition of the modular invariant of quantum tori: more precisely we

- Give an elementary definition of the (necessarily discontinuous) quantum modular invariant $j^{\text{qt}}(\theta)$ of a real number $\theta \in \mathbb{R}$, using only the distance-to-the-nearest-integer function. The association $\theta \mapsto j^{\text{qt}}(\theta)$ induces a *multi-valued* function of Mod^{qt} i.e. taking values in $2^{\mathbb{R}} \cup \{\infty\}$, which could be interpreted as the spectrum of an (as yet to be discovered) operator. Experimental evidence suggests that the set $j^{\text{qt}}(\theta)$ is finite if θ is a quadratic irrationality.
- Define a *universal* modular invariant

$$\diamond j : \diamond \widehat{\text{Mod}} \rightarrow \diamond \hat{\mathbb{C}}$$

as a *continuous* and *single valued* map of *ultrasolenoids* (see further below or §7) and show that both the classical modular invariant j^{cl} and the quantum modular invariant j^{qt} occur as subquotients of $\diamond j$. The ultrasolenoid $\diamond \widehat{\text{Mod}}$ can thus be construed as the “Riemann surface” associated to the multi-valued and discontinuous j^{qt} .

We describe in more detail how the above is accomplished by way of an overview of the sections making up this paper.

In §1 we give the definition of $j^{\text{qt}}(\theta) \in \mathbb{R} \cup \{\infty\}$, show that it is invariant with respect to the action of $\text{PGL}_2(\mathbb{Z})$ and that for $\theta \in \mathbb{Q}$, $j^{\text{qt}}(\theta) = \infty$. In §§2,3 we discuss

the case of $\theta = \varphi =$ the golden mean. We deduce an *explicit formula* for a value $j^{\text{qt}}(\varphi)$ which experiment suggests is the infimum of $j^{\text{qt}}(\varphi)$ (as well as a slight modification which appears to coincide with the experimental supremum of $j^{\text{qt}}(\varphi)$) in terms of weighted variants of the Rogers-Ramanujan functions. This concludes the part of the paper corresponding to A. above, the «elementary» part of the paper.

The remaining sections are devoted to removing the discontinuity and multivaluedness of j^{qt} by extending and lifting it to a larger space modeled on the Anosov foliation. The way that this is accomplished has the added benefit of putting the invariant into a universal geometric context, allowing us to argue in favor of its interpretation as the modular invariant of quantum tori, as well as relate it to the classical modular invariant.

In §4 we define the *generalized Kronecker foliation* $\mathcal{F}(\mu, \theta)$ of the elliptic curve $\mathbb{T}(\mu)$ by lines of μ -slope θ and establish its relation to the quantum torus $\mathbb{T}(\theta)$. We prove that the moduli space of generalized Kronecker foliations is the «Anosov foliation»

$$\text{Mod}^{\text{kf}} \approx \text{PGL}_2(\mathbb{Z}) \backslash (\pm\mathbb{H} \times (\mathbb{R} \cup \{\infty\})),$$

where $\pm\mathbb{H} = \mathbb{H} \cup \overline{\mathbb{H}}$. The leaf space of Mod^{kf} is the completed moduli space of quantum tori $\overline{\text{Mod}}^{\text{qt}} = \text{Mod}^{\text{qt}} \cup \{\infty\}$.

In §5 we review the notions of ultrafilter and ultrapower, then define (as ultrapowers over \mathbb{N}) the nonstandard integers, reals and complexes ${}^*\mathbb{Z} \subset {}^*\mathbb{R} \subset {}^*\mathbb{C}$. In §6 we define a «uniformizing lattice» for the Kronecker foliation $\mathcal{F}(\mu, \theta)$: the *diophantine approximation group*

$${}^*\Lambda(\mu, \theta) \subset {}^*\Lambda(\mu)$$

where $\Lambda(\mu) = \langle 1, \mu \rangle$ and ${}^*\Lambda(\mu)$ is its ultrapower. If ${}^*\mathbb{R} \subset {}^*\mathbb{C}$ denote the vector spaces ${}^*\mathbb{R} \subset {}^*\mathbb{C}$ modulo infinitesimals then ${}^*\Lambda(\mu, \theta) \subset {}^*\mathbb{C}$ stabilizes the line ${}^*\mathbb{R} \cdot (1 + \mu\theta)$ with quotient isomorphic to $\mathcal{F}(\mu, \theta)$.

The path is then clear: define Eisenstein series following the usual prescription – but using ${}^*\Lambda(\mu, \theta)$ in place of $\Lambda(\mu)$ – to arrive at a modular invariant that will be a (necessarily *transversally* discontinuous) function of Mod^{kf} . The challenge is to make sense of summation over the uncountable group ${}^*\Lambda(\mu, \theta)$, which we do by passing to a sheaf of ultrapowers of ${}^*\mathbb{C}$. The net effect will be an expansion of the domain and codomain of our invariant.

In fact, it makes sense to carry out this task in a universal way which shepherds both the classical and quantum invariant into the confines of a single invariant. Given $\mu \in \pm\mathbb{H}$ and $[F_\alpha] \subset {}^*\mathbb{Z}^2 - \{0, 0\}$ a *hyperfinite subset*, we may form the *hyperfinite partial sum* (see §§7,8)

$$G_k(\mu)_{[F_i]} = \sum_{({}^*m, {}^*n) \in [F_i]} ({}^*m\mu + {}^*n)^{-2k} \in {}^*\mathbb{C}.$$

One obtains a net of partial sums indexed by the set \mathcal{H} of hyperfinite subsets of ${}^*\mathbb{Z}^2 - \{0, 0\}$. This net defines a section ${}^\diamond\check{G}_k(\mu)$ of the sheaf ${}^\diamond\check{\mathbb{C}}$ of ultrapowers of ${}^*\mathbb{C}$ over the Stone space $\text{Ult}(\mathcal{H})$ of ultrafilters on \mathcal{H} . That is, evaluating the usual formula for the modular invariant at the above-defined Eisenstein sections and at the appropriate values of k , we obtain a function

$${}^\diamond\check{j}: \pm\mathbb{H} \rightarrow {}^\diamond\check{\Gamma} = \text{the set of sections of } {}^\diamond\check{\mathbb{C}}.$$

The function ${}^\diamond\check{j}$ is however *not* $\text{GL}_2(\mathbb{Z})$ -invariant, since there is an attendant shift of indices when acting by elements of $\text{GL}_2(\mathbb{Z})$. We can nevertheless achieve modularity by quotienting ${}^\diamond\check{\mathbb{C}}$ by the shift action of $\text{GL}_2(\mathbb{Z})$ on ${}^\diamond\check{\mathbb{C}}$, producing the *ultrasolenoid*

$\diamond\hat{\mathbb{C}}$. Then if we denote by $\diamond\hat{\Gamma} = \mathrm{GL}_2(\mathbb{Z}) \backslash \diamond\check{\Gamma} =$ the \mathbb{C} -algebra of *ultratransversals*, we obtain a $\mathrm{GL}_2(\mathbb{Z})$ invariant function

$$\diamond j : \pm\mathbb{H} \longrightarrow \diamond\hat{\Gamma}.$$

This is discussed in the first part of §9.

More insightfully, if we let $\diamond\mathrm{Mod}$ be the solenoid obtained as the quotient of the product $\pm\mathbb{H} \times \mathrm{Ult}(\mathcal{A}) \subset \diamond\check{\mathbb{C}}$ by the diagonal action of $\mathrm{GL}_2(\mathbb{Z})$ (shift on the base, linear action along the stalks), then we obtain, equivalently, a universal leaf preserving, transversally *continuous* function

$$\diamond j : \diamond\mathrm{Mod} \longrightarrow \diamond\hat{\mathbb{C}}.$$

The space $\diamond\mathrm{Mod}$ is an obvious generalization of the Anosov foliation $\mathrm{Mod}^{\mathrm{kf}}$, where $\mathbb{R} \cup \{\infty\}$ has been exploded and retopologized to the locally Cantor $\mathrm{Ult}(\mathcal{A})$. See Theorem 9 at the end of §9.

To recover the classical and quantum invariants, we select out ultrafilters that «observe» the groups ${}^*\Lambda(\mu)$ resp. ${}^*\Lambda(\mu, \theta)$. These are the $\mathrm{GL}_2(\mathbb{Z})$ -invariant subspaces

$$\mathrm{Cone}^{\mathrm{cl}}, \mathrm{Cone}^{\mathrm{qt}} = \bigsqcup_{\theta \in \mathbb{R}} \mathrm{Cone}^{\mathrm{qt}}(\theta) \subset \mathrm{Ult}(\mathcal{A})$$

of classical and quantum *cone ultrafilters*, see §7. They give rise to subultrasolenoids

$$\diamond\mathrm{Mod}^{\mathrm{cl}}, \diamond\mathrm{Mod}^{\mathrm{qt}} \subset \diamond\mathrm{Mod},$$

and the restriction of $\diamond j$ to each defines the classical resp. quantum invariants $\diamond j^{\mathrm{cl}}, \diamond j^{\mathrm{qt}}$. If we denote by \simeq the relation of infinitesimality, then

$$\diamond j^{\mathrm{cl}}(\mu, u) \simeq j^{\mathrm{cl}}(\mu) \quad \forall \mu \in \pm\mathbb{H}$$

moreover, every limit point $a \in j^{\mathrm{qt}}(\theta)$ is near standard to $\diamond j^{\mathrm{qt}}(i, u)$ for some $u \in \mathrm{Cone}^{\mathrm{qt}}(\theta)$. These statements are proved (at the level of ultratransversal valued invariants) in Corollary 2 and Theorem 8 of §9; their rendering into the language of ultrasolenoid valued invariants is made using Theorem 9.

In the Appendix we have presented some PARI/GP calculations which suggest that at the quadratic irrationalities, $j(\theta)$ is a finite set.

Ours is not the first attempt to use nonstandard constructions in the consideration of the RM problem: the reader may wish to compare the ideas in this paper with the work of Fesenko [9], [10] and his student Taylor [38], [39], [40]. Approaches using noncommutative geometry are discussed in the review [26].

Acknowledgments: We would like to thank B. Zil'ber for having suggested that our ideas might be applicable to the problem of RM and R. Kossak for having brought to our attention [24]. In addition, we are indebted to D. Zagier for pointing out the multivalued nature of j^{qt} and making useful suggestions for its experimental analysis. The second author would like to express his gratitude to the Isaac Newton Institute for making possible a visit in April of 2013: without which his conversations with D. Zagier would not have taken place. This work was supported in part by the grants CONACyT 058537 and PAPIIT IN103708.

1. THE QUANTUM MODULAR INVARIANT OF A REAL NUMBER

Fix $\theta \in \mathbb{R}$. Let $\|\cdot\| : \mathbb{R} \rightarrow [0, 1/2]$ denote the function which assigns to a real number its distance to the nearest integer. If $n \in \mathbb{Z}$ satisfies $\|n\theta\| < 1/2$ we denote by n^\perp the

unique closest integer, so that if we write

$$\varepsilon(n) := n\theta - n^\perp$$

then $|\varepsilon(n)| = \|n\theta\|$.

For $\varepsilon > 0$ let

$$B_\varepsilon(\theta) = \{n \in \mathbb{N} \mid \|n\theta\| < \varepsilon\}$$

and define the ε **zeta function** of θ as

$$\zeta_{\theta, \varepsilon}(s) := \sum_{B_\varepsilon(\theta)} n^{-s}.$$

Define

$$J_\varepsilon(\theta) := \frac{49}{40} \frac{\zeta_{\theta, \varepsilon}(6)^2}{\zeta_{\theta, \varepsilon}(4)^3}$$

and

$$(1) \quad j_\varepsilon(\theta) := \frac{12^3}{1 - J_\varepsilon(\theta)}.$$

Experiment indicates that the limit of $j_\varepsilon(\theta)$ does not exist as $\varepsilon \rightarrow 0$ (except for $\theta \in \mathbb{Q}$, see Proposition 1 below), but instead gives rise to a set of limit points. We indicate this state of affairs by writing

$$j^{\text{qt}}(\theta) := \lim\text{-pnt}_{\varepsilon \rightarrow 0} j_\varepsilon(\theta) \subset \mathbb{R} \cup \{\infty\}$$

where by $\lim\text{-pnt}_{\varepsilon \rightarrow 0}$ we mean the set of all limits of convergent sequences $\{j_{\varepsilon_i}(\theta)\}$, $i = 1, 2, \dots$, where $\varepsilon_i \rightarrow 0$. In this way we obtain a *multivalued map*

$$j^{\text{qt}} : \mathbb{R} \multimap \mathbb{R} \cup \{\infty\},$$

whose values may be thought of as being the spectrum of some linear operator.

There are two privileged limit points:

$$(1) \quad j^{\text{qt}}(\theta) = \liminf_{\varepsilon \rightarrow 0} j_\varepsilon^{\text{qt}}(\theta).$$

$$(2) \quad \bar{j}^{\text{qt}}(\theta) = \limsup_{\varepsilon \rightarrow 0} j_\varepsilon^{\text{qt}}(\theta).$$

There is also a privileged submultimap. Consider the sequence $\{N_i\} \subset \mathbb{N}$ of best approximations [2] to θ and define $\{\varepsilon_i^{\text{best}}\}$ by

$$\|N_i\theta\| = \varepsilon_i^{\text{best}}.$$

Then we define

$$j_{\text{best}}^{\text{qt}} : \mathbb{R} \multimap \mathbb{R} \cup \{\infty\}$$

to be the set of limit points of $\{j_{\varepsilon_i^{\text{best}}}^{\text{qt}}(\theta)\}$. We will see in the sequel that in the case of $\theta = \varphi$ the golden mean, $j_{\text{best}}^{\text{qt}}(\varphi)$ is a singleton, and experimentally satisfies (see the Appendix)

$$j_{\text{best}}^{\text{qt}}(\varphi) = \underline{j}^{\text{qt}}(\varphi).$$

Recall that the projective general linear group $\text{PGL}_2(\mathbb{Z})$ acts on $\mathbb{R} - \mathbb{Q}$ by Möbius transformations.

Theorem 1. j^{qt} and $j_{\text{best}}^{\text{qt}}$ are modular invariants:

$$j^{\text{qt}}(A(\theta)) = j^{\text{qt}}(\theta) \quad \text{and} \quad j_{\text{best}}^{\text{qt}}(A(\theta)) = j_{\text{best}}^{\text{qt}}(\theta)$$

for all $A \in \text{PGL}_2(\mathbb{Z})$.

Proof. Let

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

be a (representative of an) element of $\mathrm{PGL}_2(\mathbb{Z})$. We claim that for $\varepsilon > 0$ sufficiently small the map

$$n \mapsto cn^\perp + dn$$

defines a bijection $B_\varepsilon(\theta) \leftrightarrow B_{\varepsilon'}(A(\theta))$ where $\varepsilon' = \varepsilon \cdot |c\theta + d|^{-1}$. Indeed

$$\begin{aligned} A(\theta)(cn^\perp + dn) &= \frac{a\theta + b}{c\theta + d} \cdot (n(c\theta + d) - c\varepsilon(n)) \\ &= a\theta n + bn - c\varepsilon(n) \cdot \frac{a\theta + b}{c\theta + d} \\ &= an^\perp + bn + a\varepsilon(n) - c\varepsilon(n) \cdot \frac{a\theta + b}{c\theta + d} \\ &= an^\perp + bn + \frac{\varepsilon(n)}{c\theta + d}. \end{aligned}$$

Therefore, if $\{\varepsilon_i\}$ produces a limit point of $j^{\mathrm{qt}}(\theta)$, we have

$$\begin{aligned} \lim_{i \rightarrow \infty} J_{\varepsilon_i}(A(\theta)) &= \lim_{i \rightarrow \infty} \frac{49 \left(\sum_{n \in B_{\varepsilon_i}(\theta)} (cn^\perp + dn)^{-6} \right)^2}{40 \left(\sum_{n \in B_{\varepsilon_i}(\theta)} (cn^\perp + dn)^{-4} \right)^3} \\ &= \lim_{i \rightarrow \infty} \frac{49 \left(\sum_{n \in B_{\varepsilon_i}(\theta)} n^{-6} (c(n^\perp/n) + d)^{-6} \right)^2}{40 \left(\sum_{n \in B_{\varepsilon_i}(\theta)} n^{-4} (c(n^\perp/n) + d)^{-4} \right)^3} \\ &= \lim_{i \rightarrow \infty} \frac{49 \left(\sum_{n \in B_{\varepsilon_i}(\theta)} n^{-6} \right)^2}{40 \left(\sum_{n \in B_{\varepsilon_i}(\theta)} n^{-4} \right)^3} = \lim_{i \rightarrow \infty} J_{\varepsilon_i}(\theta), \end{aligned}$$

giving the modularity of j^{qt} . Since $A \in \mathrm{GL}_2(\mathbb{Z})$ takes tails of best approximations to tails of best approximations (see [2], page 9), a similar argument gives the modularity of $j_{\mathrm{best}}^{\mathrm{qt}}$. \square

We call $j^{\mathrm{qt}}(\theta)$ the **quantum modular invariant** of θ ; by the above result, j^{qt} defines a non continuous multivalued function

$$j^{\mathrm{qt}} : \mathrm{PGL}_2(\mathbb{Z}) \setminus \overline{\mathbb{R}} \dashrightarrow \overline{\mathbb{R}}$$

where $\overline{\mathbb{R}} = \mathbb{R} \cup \{\infty\}$.

Proposition 1. $j^{\mathrm{qt}}(\theta) = \infty$ for all $\theta \in \mathbb{Q}$.

Proof. If $\theta = q = a/b$ written in lowest terms then for ε sufficiently small, $B_\varepsilon(q) = (b)$. For such ε ,

$$J_\varepsilon(q) = \frac{49 \left(\sum_{n \in (b), n > 0} n^{-6} \right)^2}{40 \left(\sum_{n \in (b), n > 0} n^{-4} \right)^3} = \frac{49 \zeta(6)^2}{40 \zeta(4)^3} = 1$$

where the last equality follows from the Euler identities $\zeta(4) = \pi^4/2 \cdot 3^2 \cdot 5$ and $\zeta(6) = \pi^6/3^3 \cdot 5 \cdot 7$. \square

That $j^{\text{qt}} \equiv \infty$ on the rationals is in keeping with the fact that the orbit of \mathbb{Q} is regarded as the ideal boundary of the moduli space of quantum tori

$$\text{Mod}^{\text{qt}} = \text{PGL}_2(\mathbb{Z}) \backslash (\mathbb{R} - \mathbb{Q}).$$

The next two sections are devoted to calculating $j_{\text{best}}^{\text{qt}}$ of the golden mean, showing that it has, in particular, a single finite value.

2. MODULAR INVARIANT OF THE GOLDEN MEAN I: AN EXPLICIT FORMULA

Let

$$\varphi := \frac{1 + \sqrt{5}}{2}$$

be the golden mean. In this section we will produce, assuming that $j_{\text{best}}^{\text{qt}}(\varphi)$ converges, an explicit formula for $j_{\text{best}}^{\text{qt}}(\varphi)$ obtained by evaluating at φ a certain rational expression involving weighted variants of the Rogers-Ramanujan functions. The convergence of $j_{\text{best}}^{\text{qt}}(\varphi)$ will be proved in §3; in the Appendix we will present evidence that suggests that its value is the minimum of $j^{\text{qt}}(\varphi)$.

We begin by recalling some facts about the golden mean and its diophantine approximations, see for example [33], [41]. The minimal polynomial of φ is $X^2 - X - 1$ and φ is a unit in $\mathbb{Q}(\sqrt{5})$, whose inverse is -1 times its conjugate:

$$\varphi^{-1} = -\varphi' = \frac{\sqrt{5} - 1}{2}.$$

The discriminant of φ is $\sqrt{5}$, and the class number of $\mathbb{Q}(\sqrt{5})$ is one. The pseudo lattice $\langle 1, \varphi \rangle$ has endomorphism ring equal to O_K , hence has conductor $f = 1$.

If we denote by $[a_0, a_1, \dots]$ the sequence of partial quotients of a real number θ then for $\theta = \varphi$, $a_i = 1$ for all i . It follows that the sequence of best approximations (p_m, q_m) of φ is given by (F_{m+1}, F_m) , where $\{F_m\} = \{1, 1, 2, 3, 5, 8, \dots\}$, $m \geq 1$, denotes the Fibonacci sequence:

$$F_{m+1} = F_m + F_{m-1}, \quad m \geq 1.$$

See for example [33]. This means that as $m \rightarrow \infty$,

$$(2) \quad \varepsilon_m := F_m \varphi - F_{m+1} \longrightarrow 0$$

and that for all $0 < n < F_m$,

$$\|n\varphi\| > \|F_m \varphi\| = |\varepsilon_m|,$$

where as before $\|x\|$ is the distance of x to the nearest integer.

We recall Binet's formula [29]:

$$F_m = \frac{\varphi^m - (\varphi')^m}{\sqrt{5}} = \frac{\varphi^m - (-1)^m \varphi^{-m}}{\sqrt{5}} = \begin{cases} \frac{\varphi^m - \varphi^{-m}}{\sqrt{5}} & \text{if } m \text{ is even} \\ \frac{\varphi^m + \varphi^{-m}}{\sqrt{5}} & \text{if } m \text{ is odd.} \end{cases}$$

Using Binet's formula, we may obtain the following explicit expression for ε_m of (2):

$$(3) \quad \varepsilon_m = (-1)^{m+1} \varphi^{-m}.$$

Indeed, for each integer m we have

$$\begin{aligned} F_m \varphi - F_{m+1} &= \left(\frac{\varphi^m + (-1)^{m+1} \varphi^{-m}}{\sqrt{5}} \right) \varphi - \left(\frac{\varphi^{m+1} + (-1)^m \varphi^{-m-1}}{\sqrt{5}} \right) \\ &= \frac{1}{\sqrt{5}} (\varphi^{m+1} + (-1)^{m+1} \varphi^{-m+1} - \varphi^{m+1} + (-1)^{m+1} \varphi^{-m-1}) \\ &= \frac{1}{\sqrt{5}} (\varphi + \varphi^{-1}) (-1)^{m+1} \varphi^{-m} = (-1)^{m+1} \varphi^{-m}. \end{aligned}$$

Notice then that for $m \geq 2$, we have

$$\|F_m \varphi\| = |\varepsilon_m|$$

and in particular,

$$\varepsilon_m^{\text{best}} = \varphi^{-m}.$$

For m large, $\sqrt{5}F_m \approx \varphi^m$, with an error term $= \pm \varphi^{-m}$ that decays exponentially as $m \rightarrow \infty$.

Finally, we recall Zeckendorf's representation (which is actually a special case of a more general result of Ostrowski [30]):

Theorem 2 (Zeckendorf, [42]). *Every natural number $n \in \mathbb{N}$ may be written uniquely as a sum of non-consecutive Fibonacci numbers:*

$$n = F_I := F_{i_1} + \dots + F_{i_k}, \quad 2 \leq i_1, i_1 + 2 \leq i_2, \dots, i_{k-1} + 2 \leq i_k, \quad 1 \leq k.$$

Note 1. The condition that $i_1 \geq 2$ is to ensure uniqueness in the decomposition, otherwise the value 1 could occur in two different ways, as F_1 or F_2 .

We now develop an explicit formula for $J_{\text{best}}^{\text{qt}}(\varphi)$. Write $\varepsilon = |\varepsilon_m|$ and $B = B_m(\varphi) = \{n \in \mathbb{N} \mid \|n\varphi\| < \varepsilon\}$ so that

$$J_{\varepsilon}^{\text{qt}}(\varphi) = \frac{49}{40} \frac{(\sum_{n \in B} n^{-6})^2}{(\sum_{n \in B} n^{-4})^3}.$$

The first step is to determine the elements of B in terms of their Zeckendorf representations. In what follows, for a multi-index $I = (i_1, \dots, i_k)$, define $|I| = k$.

Lemma 1. *Let $n = F_I = F_{i_1} + \dots + F_{i_k}$ written in its unique Zeckendorf form. Then $n \in B$ if and only*

- I. $|I| \geq 1$, $i_1 \geq m + 1$ or
- II. $|I| \geq 2$, $i_1 = m$ and $i_2 - m$ is odd.

Note 2. Since the Zeckendorf form consists of sums of nonconsecutive Fibonacci numbers, we must have that $i_2 - m \geq 3$ in II.

Proof. First note that we have trivially by (3) that $F_{m+i} \in B$ for $i \geq 1$. Suppose that $n = F_I$ is a sum of more than one non-consecutive Fibonacci numbers and $i_1 \geq m + 1$. Then we have

$$\|n\varphi\| < \varphi^{-(m+1)} + \varphi^{-(m+3)} + \dots = \varphi^{-(m+1)}(1 - \varphi^{-2})^{-1}.$$

Since $\varphi = \varphi^2 - 1$ it follows that $(1 - \varphi^{-2})^{-1} = \varphi$. Then $\|n\varphi\| < \varphi^{-m}$ which implies that $n \in B$. Thus every element of the type described in I. belongs to B . On the other hand, if $i_1 \leq m - 1$, then we claim that

$$\varphi^{-m} = \varepsilon < \|n\varphi\| < 1 - \varphi^{-1} = \varphi^{-2}.$$

Indeed, if $n = F_I$, the associated error term sum

$$\varepsilon_I := \pm \varepsilon_{i_1} \pm \cdots \pm \varepsilon_{i_k}$$

is minimized in absolute value by taking $i_1 = m - 1$ and assuming that the remaining indices i_2, \dots are such that the signs of the associated error terms ε_{i_2}, \dots are different from the sign of the error term ε_{m-1} . More precisely,

$$|\varepsilon_I| > \varphi^{-(m-1)} - (\varphi^{-(m+2)} + \varphi^{-(m+4)} + \cdots) = \varphi^{-m}(\varphi - \varphi^{-2}(1 - \varphi^{-2})^{-1}).$$

Since $\varphi^{-2}(1 - \varphi^{-2})^{-1} = \varphi^{-1}$ and $\varphi - \varphi^{-1} = 1$, it follows that $|\varepsilon_I| > \varphi^{-m} = \varepsilon$. In addition $|\varepsilon_I|$ is maximized by taking $i_1 = 2, i_2 = 4, \dots$, so that

$$|\varepsilon_I| < \varphi^{-2} + \varphi^{-4} + \cdots = \frac{1}{\varphi^2 - 1} = \varphi^{-1}.$$

Note that the distance of the latter bound φ^{-1} to the nearest integer is $1 - \varphi^{-1} = \varphi^{-2}$. It follows then from the definition of $\|\cdot\|$ and the fact that we are assuming that $m > 2$ that $\|n\varphi\| > \varphi^{-m} = \varepsilon$ and $n \notin B$. Now if $i_1 = m$ and $i_2 - m$ is even, then the error terms ε_m and ε_{i_2} share the same sign, and we have

$$\|n\varphi\| > \varphi^{-m} + \varphi^{-i_2} - (\varphi^{-(i_2+3)} + \varphi^{-(i_2+5)} + \cdots) = \varepsilon + (\varphi^{-i_2} - \varphi^{-(i_2+3)}(1 - \varphi^{-2})^{-1}) > \varepsilon$$

Indeed, the last inequality follows since

$$\begin{aligned} & \varphi^{-i_2} - \varphi^{-(i_2+3)}(1 - \varphi^{-2})^{-1} = \\ & \varphi^{-i_2}(1 - \varphi^{-3}(1 - \varphi^{-2})^{-1}) = \\ & \varphi^{-i_2}(1 - \varphi^{-2}(\varphi - \varphi^{-1})^{-1}) = \\ & \varphi^{-i_2}(1 - \varphi^{-2}) > 0. \end{aligned}$$

On the other hand, if $i_1 = m$ and $i_2 = m + k$, k odd, then the sign of the corresponding error terms differ, and we have

$$\begin{aligned} \|n\varphi\| & < \varphi^{-m} - \varphi^{-m-k} + \varphi^{-m-k-3} + \varphi^{-m-k-5} + \cdots \\ & = \varphi^{-m} - \varphi^{-m-k}(1 - (\varphi^{-3} + \varphi^{-5} + \cdots)) \\ & = \varphi^{-m} - \varphi^{-m-k}(1 - \varphi^{-3}(1 - \varphi^{-2})^{-1}) \\ & = \varphi^{-m} - \varphi^{-m-k}(1 - \varphi^{-2}) < \varepsilon \end{aligned}$$

so that $n \in B$. □

Let \mathfrak{N} be the set of increasing, finite tuples $I = (i_1, \dots, i_l)$ of natural numbers with $|I| = l \geq 2$ and which are not consecutive i.e. $i_1 + 2 \leq i_2, \dots, i_{l-1} + 2 \leq i_l$. Denote by

$$(4) \quad \mathfrak{N}(m) = \{I = (i_1, \dots, i_l) \in \mathfrak{N} \mid i_1 \geq m\}.$$

Also denote by

$$(5) \quad \mathfrak{M}(m) = \{I \in \mathfrak{N}(m) \mid i_1 = m \text{ and } i_2 = m + k \text{ for } k \text{ odd}\}.$$

Consider B_m for $m > 2$. Then by the Lemma we have

$$J_{\varepsilon_m}^{\text{qt}}(\varphi) = \frac{49 \left(\sum_{n \in B_m} n^{-6} \right)^2}{40 \left(\sum_{n \in B_m} n^{-4} \right)^3} = \frac{49 \left(\sum_{i=1}^{\infty} F_{m+i}^{-6} + \sum_{I \in \mathfrak{N}(m+1)} F_I^{-6} + \sum_{I \in \mathfrak{M}(m)} F_I^{-6} \right)^2}{40 \left(\sum_{i=1}^{\infty} F_{m+i}^{-4} + \sum_{I \in \mathfrak{N}(m+1)} F_I^{-4} + \sum_{I \in \mathfrak{M}(m)} F_I^{-4} \right)^3},$$

an expression whose status is still only formal. Consider also the formal expression

$$(6) \quad J_0^{\text{qt}}(\varphi) := \frac{49 \left(\sum_{i=1}^{\infty} \varphi^{-6i} + \sum_{I \in \mathfrak{N}(1)} \varphi_I^{-6} + \sum_{I \in \mathfrak{N}(0)} \varphi_I^{-6} \right)^2}{40 \left(\sum_{i=1}^{\infty} \varphi^{-4i} + \sum_{I \in \mathfrak{N}(1)} \varphi_I^{-4} + \sum_{I \in \mathfrak{N}(0)} \varphi_I^{-4} \right)^3}$$

where

$$\varphi_I := \varphi^{i_1} + \dots + \varphi^{i_l}.$$

Theorem 3. *If $J_0^{\text{qt}}(\varphi)$ converges then so does $J_{\varepsilon_m}^{\text{qt}}(\varphi)$ for each m and*

$$J_{\varepsilon_m}^{\text{qt}}(\varphi) \longrightarrow J_0^{\text{qt}}(\varphi) = J_{\text{best}}^{\text{qt}}(\varphi)$$

as $m \rightarrow \infty$.

Proof. Multiply the numerator and denominator of $J_{\varepsilon_m}^{\text{qt}}(\varphi)$ by F_m^{12} to obtain

$$(7) \quad J_{\varepsilon_m}^{\text{qt}}(\varphi) = \frac{49 \left(\sum_{i=1}^{\infty} (F_m/F_{m+i})^6 + \sum_{I \in \mathfrak{N}(m+1)} (F_m/F_I)^6 + \sum_{I \in \mathfrak{N}(m)} (F_m/F_I)^6 \right)^2}{40 \left(\sum_{i=1}^{\infty} (F_m/F_{m+i})^4 + \sum_{I \in \mathfrak{N}(m+1)} (F_m/F_I)^4 + \sum_{I \in \mathfrak{N}(m)} (F_m/F_I)^4 \right)^3}.$$

It will suffice to show that each term $T_m^{-6} = T_{m,I}^{-6}$ ($T_m^{-4} = T_{m,I}^{-4}$) appearing in a sum contained in the numerator (denominator) of (7) satisfies

$$C_m^{-6} \cdot T^{-6} < T_m^{-6} < C_m^6 \cdot T^{-6} \quad \left(C_m^{-4} \cdot T^{-4} < T_m^{-4} < C_m^4 \cdot T^{-4} \right)$$

where $T = T_I$ is the correspondingly indexed term of $J_0^{\text{qt}}(\varphi)$ and

$$C_m = \frac{1 + \varphi^{-2m}}{1 - \varphi^{-2m}}.$$

This will give convergence of each $J_{\varepsilon_m}^{\text{qt}}(\varphi)$, as well as the bound

$$\left(\frac{1 - \varphi^{-2m}}{1 + \varphi^{-2m}} \right)^{24} J_0^{\text{qt}}(\varphi) < J_{\varepsilon_m}^{\text{qt}}(\varphi) < \left(\frac{1 + \varphi^{-2m}}{1 - \varphi^{-2m}} \right)^{24} J_0^{\text{qt}}(\varphi),$$

which implies that $J_{\varepsilon_m}^{\text{qt}}(\varphi) \rightarrow J_0^{\text{qt}}(\varphi)$.

We will now make use of Binet's formula, $\sqrt{5}F_m = (\varphi^m \pm \varphi^{-m})$. Note that the $\sqrt{5}$ factors drop out and so we may simply replace every Fibonacci term F_m appearing by $\varphi^m \pm \varphi^{-m}$.

We consider first the numerator of (7), treating each of the three sums there separately. The first sum may be written

$$\sum_{i=1}^{\infty} (F_m/F_{m+i})^6 = \sum_{i=1}^{\infty} \left(\frac{\varphi^m \pm \varphi^{-m}}{\varphi^{m+i} \pm (-1)^i \varphi^{-(m+i)}} \right)^6 = \sum_{i=1}^{\infty} \varphi^{-6i} \left(\frac{1 \pm \varphi^{-2m}}{1 \pm (-1)^i \varphi^{-2m-2i}} \right)^6.$$

Note that

$$\left(\frac{1 - \varphi^{-2m}}{1 + \varphi^{-2m}} \right)^6 < \left(\frac{1 \pm \varphi^{-2m}}{1 \pm (-1)^i \varphi^{-2m-2i}} \right)^6 < \left(\frac{1 + \varphi^{-2m}}{1 - \varphi^{-2m}} \right)^6.$$

The next sum is

$$(8) \quad \sum_{I \in \mathfrak{N}(m+1)} (F_m/F_I)^6 = \sum_{I \in \mathfrak{N}(m+1)} \left(\frac{\varphi^m \pm \varphi^{-m}}{(\varphi^{m+i_1} \pm \varphi^{-m-i_1}) + \dots + (\varphi^{m+i_k} \pm \varphi^{-m-i_k})} \right)^6,$$

where we are writing our generic $I \in \mathfrak{N}(m+1)$ in the form $I = (i_1 + m, \dots, i_k + m)$ with $1 \leq i_1 < i_2 < \dots < i_k$. Letting $I_0 = (i_1, \dots, i_k)$ then each term of the sum in (8) may be re-written

$$(9) \quad \left(\frac{1 \pm \varphi^{-2m}}{\varphi_{I_0} + (\pm \varphi_{-I_0-2m})} \right)^6 = \varphi_{I_0}^{-6} \cdot \left(\frac{1 \pm \varphi^{-2m}}{1 + (\pm \varphi_{-I_0-2m})/\varphi_{I_0}} \right)^6$$

where

$$\pm \varphi_{-I_0-2m} := \pm \varphi^{-i_1-2m} \pm \dots \pm \varphi^{-i_k-2m},$$

the signs determined as in Binet's formula by the parities of the powers. It is easy to see that

$$(10) \quad \left(\frac{1 - \varphi^{-2m}}{1 + \varphi^{-2m}} \right)^6 < \left(\frac{1 \pm \varphi^{-2m}}{1 + (\pm \varphi_{-I_0-2m})/\varphi_{I_0}} \right)^6 < \left(\frac{1 + \varphi^{-2m}}{1 - \varphi^{-2m}} \right)^6 :$$

indeed, both inequalities in (10) follow since

$$\varphi^{-2m} > (\pm \varphi_{-I_0-2m})/\varphi_{I_0} > -\varphi^{-2m},$$

true as

$$(11) \quad (\pm \varphi_{-I_0-2m})/\varphi_{I_0} = \varphi^{-2m} \left(\frac{\pm \varphi^{-i_1} \pm \dots \pm \varphi^{-i_k}}{\varphi^{i_1} + \dots + \varphi^{i_k}} \right).$$

What remains is the sum over $\mathfrak{M}(m)$: the analysis here is essentially the same as that made for the sum over $\mathfrak{N}(m+1)$, only we take into account that $I = (m, m+j, m+i_3, \dots, m+i_k)$ where j is odd. Writing $I_0 = (0, j, i_3, \dots, i_k)$, then we have the equation (9) with

$$\pm \varphi_{-I_0-2m} = \pm \varphi^{-2m} \mp \varphi^{-j-2m} \pm \dots \pm \varphi^{-i_k-2m},$$

where the \mp sign of φ^{-j-2m} indicates that this sign is opposite to that of φ^{-2m} , as j is odd. The analogue of (11) is then

$$(\pm \varphi_{-I_0-2m})/\varphi_{I_0} = \varphi^{-2m} \left(\frac{\pm 1 \mp \varphi^{-j} \pm \dots \pm \varphi^{-i_k}}{1 + \varphi^j + \dots + \varphi^{i_k}} \right),$$

which yields the analogue of (10) in this case. This completes our bounding of the numerator. Analogous bounds, with the exponent 6 replaced by 4, may be found for the corresponding sums in the denominator of $\mathcal{J}_{\varepsilon_m}^{\text{qt}}$. The result now follows. \square

Let $P(n)$ be the set of partitions of n into distinct parts whose differences are at least 2, and let $c(n) = |P(n)|$. The generating function

$$F(x) = \sum_{n=1}^{\infty} c(n)x^n = \sum \frac{x^{n^2}}{(1-x) \cdots (1-x^n)}$$

is of substantial combinatorial interest: $1 + F(x)$ is the left-hand side of the first Rogers-Ramanujan identity [17].

For each partition $I \in P(n)$, let $f_I(x) = x^{i_1} + \dots + x^{i_k}$ be the associated weighting polynomial. Define

$$C_{x,M}(n) = x^{Mn} \sum_{I \in P(n)} f_I(x)^{-M}.$$

Consider the generating function

$$G_M(x) = \sum C_{x,M}(n)x^n.$$

Clearly we have

$$G_M(\varphi) = \sum_{i=1}^{\infty} \varphi^{-Mi} + \sum_{I \in \mathfrak{N}(1)} \varphi_I^{-M}.$$

Similarly, let $\mathcal{Q}(n) \subset P(n)$ be the set of those partitions $I = i_1 < i_2 < \dots < i_k$ in $P(n)$ for which i_1 is odd and ≥ 3 . Let

$$D_{x,M}(n) := x^{Mn} \sum_{I \in \mathcal{Q}(n)} (1 + f_I(x))^{-M}$$

and define

$$H_M(x) := \sum D_{x,M}(n) x^n.$$

Then

$$H_M(\varphi) = \sum_{I \in \mathfrak{M}(0)} \varphi_I^{-M}.$$

The following is then immediate:

Corollary 1. *Let $J_0^{\text{qt}}(\varphi)$ be as above. Then*

$$(12) \quad J_0^{\text{qt}}(\varphi) = \frac{49 (G_6(\varphi) + H_6(\varphi))^2}{40 (G_4(\varphi) + H_4(\varphi))^3}.$$

Note 3. If one replaces in the formula for $C_{x,M}(n)$ the weighting polynomial $f_I(x)^{-M}$ by the equiweight x^{-Mn} one recovers $c(n)$. Thus the functions $G_M(x), H_M(x)$ may be viewed as weighted variants of the variable part of the Rogers-Ramanujan function.

In the Appendix, we show that by replacing $G_M(\varphi)$ by $G'_M(\varphi) = G_M(\varphi) + 1$ in (12) we obtain a value very close to the experimental supremum of $J^{\text{qt}}(\varphi)$.

3. MODULAR INVARIANT OF THE GOLDEN MEAN II: CONVERGENCE

In this section we will show that $j_{\text{best}}^{\text{qt}}(\varphi) < \infty$ and in fact converges. As before we write $j_{\text{best}}^{\text{qt}}(\varphi) := 12^3 / (1 - J_{\text{best}}^{\text{qt}}(\varphi))$.

Theorem 4. *$J_{\text{best}}^{\text{qt}}(\varphi)$ converges with the bounds*

$$9150 < J_{\text{best}}^{\text{qt}}(\varphi) < 9840.$$

Proof. To prove the convergence of $J_{\text{best}}^{\text{qt}}(\varphi)$, it is enough to prove convergence of the explicit formula $J_0^{\text{qt}}(\varphi) := 12^3 / (1 - J_{\text{best}}^{\text{qt}}(\varphi))$ obtained from (6). Observe first that

$$\sum_{i=1}^{\infty} \varphi^{-6i} = (\varphi^6 - 1)^{-1}, \quad \sum_{i=1}^{\infty} \varphi^{-4i} = (\varphi^4 - 1)^{-1}$$

so we may write

$$J_0^{\text{qt}}(\varphi) = \frac{49 \left((\varphi^6 - 1)^{-1} + \sum_{I \in \mathfrak{N}(1)} \varphi_I^{-6} + \sum_{I \in \mathfrak{M}(0)} \varphi_I^{-6} \right)^2}{40 \left((\varphi^4 - 1)^{-1} + \sum_{I \in \mathfrak{N}(1)} \varphi_I^{-4} + \sum_{I \in \mathfrak{M}(0)} \varphi_I^{-4} \right)^3}.$$

We now find an explicit approximation and an upper bound for the sum $\sum_{I \in \mathfrak{N}(1)} \varphi_I^{-M}$ where M is a positive integer. In fact, we will show that

$$(13) \quad \sum_{I \in \mathfrak{N}(1)} \varphi_I^{-M} = \frac{1}{(\varphi^M - 1)(\varphi^2 + 1)^M} + C(M)$$

where

$$(14) \quad C(M) < \tilde{C}(M) := \frac{1}{\varphi^{2M}(\varphi^M - 1)^2} + \frac{1}{\varphi^M(\varphi^M - 1)^2(\varphi^{2M} - \varphi^M - 1)}.$$

Consider first the sum of those I with $|I| = 2$:

$$(15) \quad \sum_{\substack{i_1 \geq 1 \\ i_2 \geq i_1+2}} \frac{1}{(\varphi^{i_1} + \varphi^{i_2})^M} = \sum_{i=1}^{\infty} \varphi^{-Mi} \sum_{k=2}^{\infty} (1 + \varphi^k)^{-M}$$

$$= \frac{1}{\varphi^M - 1} \left\{ \frac{1}{(1 + \varphi^2)^M} + \sum_{k=3}^{\infty} (1 + \varphi^k)^{-M} \right\}$$

$$(16) \quad < \frac{1}{\varphi^M - 1} \left\{ \frac{1}{(1 + \varphi^2)^M} + \sum_{k=3}^{\infty} \varphi^{-Mk} \right\}$$

$$= \frac{1}{(\varphi^M - 1)(\varphi^2 + 1)^M} + \frac{1}{\varphi^{2M}(\varphi^M - 1)^2}.$$

The equality (15) produces the explicit term $1/((\varphi^M - 1)(\varphi^2 + 1)^M)$ appearing in (13); the second term in (16) is the first bounding term in (14).

For $|I| = 3$ we have

$$\sum_{\substack{i_1 \geq 1 \\ i_2 \geq i_1+2, i_3 \geq i_2+2}} \frac{1}{(\varphi^{i_1} + \varphi^{i_2} + \varphi^{i_3})^M} = \sum_{\substack{i_1 \geq 1 \\ i_2 \geq i_1+2, i_3 \geq i_2+2}} \varphi^{-Mi_1} \frac{1}{(1 + \varphi^{i_2-i_1} + \varphi^{i_3-i_1})^M}$$

$$< \sum_{\substack{i_1 \geq 1 \\ i_2 \geq i_1+2, i_3 \geq i_2+2}} \varphi^{-Mi_1} \frac{1}{(\varphi^{i_2-i_1} + \varphi^{i_3-i_1})^M}$$

$$= \sum_{\substack{i_1 \geq 1 \\ i_2 \geq i_1+2, i_3 \geq i_2+2}} \varphi^{-Mi_1} \varphi^{-M(i_2-i_1)} \frac{1}{(1 + \varphi^{i_3-i_2})^M}$$

$$< \sum_{i=1}^{\infty} \varphi^{-Mi} \sum_{j \geq 2} \varphi^{-Mj} \sum_{k \geq 2} \varphi^{-Mk}$$

$$= \frac{(\varphi^{-M})^2}{(\varphi^M - 1)^3}.$$

Inductively, for the terms with $|I| = l \geq 3$ we have the bound

$$\frac{(\varphi^{-M})^{l-1}}{(\varphi^M - 1)^l}.$$

Summing these bounds from $l = 3$ to ∞ gives the second term in (14):

$$\sum_{l=3}^{\infty} \frac{(\varphi^{-M})^{l-1}}{(\varphi^M - 1)^l} = \varphi^M \sum_{l=3}^{\infty} \frac{1}{(\varphi^M(\varphi^M - 1))^l} = \frac{1}{\varphi^M(\varphi^M - 1)^2(\varphi^{2M} - \varphi^M - 1)}$$

We now bound the second type of sum appearing in $\mathcal{J}_0^{\text{qt}}(\varphi)$, $\sum_{I \in \mathfrak{M}(0)} \varphi_I^{-M}$. We will show here that

$$(17) \quad \sum_{I \in \mathfrak{M}(0)} \varphi_I^{-M} = \frac{1}{(1 + \varphi^3)^M} + D(M)$$

where

$$(18) \quad D(M) < \tilde{D}(M) := \frac{1}{\varphi^{3M}(\varphi^{2M} - 1)} + \frac{1}{\varphi^M(\varphi^{2M} - 1)(\varphi^{2M} - \varphi^M - 1)}$$

When $|I| = 2$ we have, since $i_1 = 0$, that $i_2 = 2j + 1$ is odd, where $j \geq 1$ (recall the definition of $\mathfrak{M}(m)$ found in (5)). For such I we have the contribution

$$\begin{aligned}
(19) \quad \sum_{\substack{i=2j+1 \\ j \geq 1}} \frac{1}{(1 + \varphi^i)^M} &= \frac{1}{(1 + \varphi^3)^M} + \sum_{j=2}^{\infty} (1 + \varphi^{(2j+1)})^{-M} \\
&< \frac{1}{(1 + \varphi^3)^M} + \sum_{j=2}^{\infty} \varphi^{-M(2j+1)} \\
&= \frac{1}{(1 + \varphi^3)^M} + \varphi^{-M} \sum_{j=2}^{\infty} \varphi^{-2Mj} \\
&= \frac{1}{(1 + \varphi^3)^M} + \varphi^{-5M} \frac{1}{1 - \varphi^{-2M}} \\
(20) \quad &= \frac{1}{(1 + \varphi^3)^M} + \frac{1}{\varphi^{3M}(\varphi^{2M} - 1)}.
\end{aligned}$$

For $|I| = 3$ we have

$$\begin{aligned}
\sum_{j \geq 1, k \geq (2j+1)+2} \frac{1}{(1 + \varphi^{2j+1} + \varphi^k)^M} &< \sum_{j \geq 1, k \geq (2j+1)+2} \varphi^{-M(2j+1)} \frac{1}{(1 + \varphi^{k-(2j+1)})^M} \\
&< \sum_{j=1}^{\infty} \varphi^{-M(2j+1)} \sum_{k=2}^{\infty} \varphi^{-Mk} \\
&= \frac{1}{\varphi^M(\varphi^{2M} - 1)} \cdot \frac{1}{\varphi^M(\varphi^M - 1)} \\
&= \frac{1}{\varphi^M + 1} \cdot \left(\frac{\varphi^{-M}}{\varphi^M - 1} \right)^2
\end{aligned}$$

For the sum over I with $|I| = l$, we obtain inductively the bound

$$\frac{1}{\varphi^M + 1} \left(\frac{\varphi^{-M}}{\varphi^M - 1} \right)^{l-1}$$

and summing these from $l = 3$ to ∞ gives

$$\frac{1}{\varphi^M(\varphi^{2M} - 1)(\varphi^{2M} - \varphi^M - 1)}.$$

It follows then that

$$\mathcal{J}_0^{\text{qt}}(\varphi) < \frac{49}{40} \frac{\left((\varphi^6 - 1)^{-1} + ((\varphi^6 - 1)(\varphi^2 + 1)^6)^{-1} + (1 + \varphi^3)^{-6} + \tilde{C}(6) + \tilde{D}(6) \right)^2}{\left((\varphi^4 - 1)^{-1} + ((\varphi^4 - 1)(\varphi^2 + 1)^4)^{-1} + (1 + \varphi^3)^{-4} \right)^3}$$

$$\approx 0.824376700276.$$

A lower bound may be given by

$$0.81115979990388 \approx \frac{49}{40} \frac{\left((\varphi^6 - 1)^{-1} + ((\varphi^6 - 1)(\varphi^2 + 1)^6)^{-1} + (1 + \varphi^3)^{-6} \right)^2}{\left((\varphi^4 - 1)^{-1} + ((\varphi^4 - 1)(\varphi^2 + 1)^4)^{-1} + (1 + \varphi^3)^{-4} + \tilde{C}(4) + \tilde{D}(4) \right)^3}$$

$$< \mathcal{J}_0^{\text{qt}}(\varphi)$$

which give the bounds presented in the statement of the theorem. Since the numerator and denominator of $J_0^{\text{qt}}(\varphi)$ are hypergeometric functions with positive coefficients evaluated at a positive real number, it follows from the above bounds that they converge, and in particular, that $J_0^{\text{qt}}(\varphi)$ converges. \square

Note 4. Using the PARI/GP value of the explicit formula of Corollary 1, we get $j_{\text{best}}^{\text{qt}}(\varphi) \approx 9538.249655644$, which agrees closely with the experimental value obtained for $\underline{j}^{\text{qt}}(\varphi)$. See the Appendix.

4. QUANTUM TORI AND KRONECKER FOLIATIONS

In this section we begin the process of finding a continuous and single valued version of the set-valued quantum invariant j^{qt} , as well as putting the latter in its proper geometrical context.

Consider the Kronecker foliation $\mathcal{F}(\theta)$ of slope θ in the elliptic curve $\mathbb{T}(i) = \mathbb{C}/\langle 1, i \rangle$, i.e. the image in $\mathbb{T}(i)$ of the foliation of the complex plane \mathbb{C} by lines of slope θ . The leaf space of $\mathcal{F}(\theta)$ may be identified with the quotient group $\mathbb{T}(i)/L(\theta)$, where $L(\theta)$ is the leaf through the origin, a 1-parameter subgroup of $\mathbb{T}(i)$. When $\theta \in \mathbb{R} - \mathbb{Q}$, $L(\theta)$ is dense in $\mathbb{T}(i)$ so that the leaf space is non Hausdorff.

On the other hand, let $\Lambda(\theta) = \langle 1, \theta \rangle \subset \mathbb{R}$ be the pseudo lattice generated by 1 and θ . As discussed in the Introduction, the quantum torus associated to $\theta \in \mathbb{R}$ may be defined as the following quotient:

$$\mathbb{T}(\theta) = \mathbb{R}/\Lambda(\theta).$$

When θ is irrational, this is a non Hausdorff topological group. It will be convenient for us to allow θ to be rational as well, in which case one obtains the circle.

Proposition 2. *The leaf space of $\mathcal{F}(\theta)$ is canonically isomorphic to $\mathbb{T}(\theta)$.*

Proof. Writing $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$, consider the suspension $(\mathbb{R} \times \mathbb{S}^1)/\mathbb{Z}$, where the action of \mathbb{Z} is diagonal: $n \cdot (r, s + \mathbb{Z}) = (r + n, (s - \theta n) + \mathbb{Z})$. The suspension defines a linear foliation of $\mathbb{T}(i)$: the image of the product foliation $\mathbb{R} \times \mathbb{S}^1$, whose leaves are of the form $\mathbb{R} \times \{s + \mathbb{Z}\}$. There is an isomorphism of the Kronecker foliation $\mathcal{F}(\theta)$ with this foliation, induced by $\mathbb{C} \rightarrow (\mathbb{R} \times \mathbb{S}^1)$, $r + is \mapsto (r, s - r\theta + \mathbb{Z})$. Through this identification, one sees that the leaf space of $\mathcal{F}(\theta)$ is canonically isomorphic to the quotient group $\mathbb{S}^1/\langle \theta + \mathbb{Z} \rangle$. But the latter is canonically isomorphic to $\mathbb{T}(\theta)$. \square

The Kronecker foliation has an obvious generalization in which one replaces $\mathbb{T}(i)$ by any elliptic curve $\mathbb{T}(\mu) = \mathbb{C}/\Lambda(\mu)$ where $\Lambda(\mu) = \langle 1, \mu \rangle$ and where $\mu \in \mathbb{H} =$ the hyperbolic plane. Given $\theta \in \mathbb{R} \cup \{\infty\} \approx \mathbb{S}^1$, let $\widetilde{\mathcal{F}}(\mu, \theta)$ be the foliation of \mathbb{C} defined by the translates of the line of « μ -slope θ »,

$$\widetilde{L}(\mu, \theta) = \begin{cases} \mathbb{R} \cdot (1 + \theta\mu) & \text{if } \theta \neq \infty \\ \mathbb{R} \cdot \mu & \text{if } \theta = \infty \end{cases}$$

The image $\mathcal{F}(\mu, \theta)$ of $\widetilde{\mathcal{F}}(\mu, \theta)$ in $\mathbb{T}(\mu)$ is called a **generalized Kronecker foliation** of slope θ and modulus μ . Alternatively, $\mathcal{F}(\mu, \theta)$ is completely determined by the pair

$$(\mathbb{T}(\mu), L(\mu, \theta))$$

consisting of the elliptic curve and the distinguished 1-parameter subgroup

$$L(\mu, \theta) = \text{image of the line } \widetilde{L}(\mu, \theta) = \text{leaf through } 0.$$

This may be regarded as a continuous generalization of the notion of an elliptic curve equipped with a distinguished finite subgroup of order N .

As in [25], it will be convenient to allow the parameter μ to take values in $\overline{\mathbb{H}}$ as well. If we denote by $\pm\mathbb{H} = \mathbb{H} \cup \overline{\mathbb{H}}$ then $\mathrm{PGL}_2(\mathbb{Z})$ acts on $\pm\mathbb{H}$ by isometries and we recover by quotient the classical moduli space of elliptic curves

$$\mathrm{Mod}^{\mathrm{cl}} := \mathrm{PSL}_2(\mathbb{Z}) \backslash \mathbb{H} \approx \mathrm{PGL}_2(\mathbb{Z}) \backslash \pm\mathbb{H}.$$

The Kronecker foliation $\mathcal{F}(\mu, \theta)$ for $\mu \in \overline{\mathbb{H}}$ is defined exactly as in the case of $\mu \in \mathbb{H}$. Note that for all $(\mu, \theta) \in \pm\mathbb{H} \times \mathbb{S}^1$ we have the equality

$$\mathcal{F}(-\mu, -\theta) = \mathcal{F}(\mu, \theta).$$

This equality remains true for $\theta = \infty$ (which, like 0, has no sign).

Let $(\mu, \theta), (\mu', \theta') \in \pm\mathbb{H} \times \mathbb{S}^1$. The Kronecker foliations $\mathcal{F}(\mu, \theta)$ and $\mathcal{F}(\mu', \theta')$ are said to be equivalent if there exists a homothety $z \mapsto \lambda z$ inducing an isomorphism of underlying elliptic curves that transports $\mathcal{F}(\mu, \theta)$ to $\mathcal{F}(\mu', \theta')$: or equivalently, inducing an isomorphism of pairs

$$f : (\mathbb{T}(\mu), L(\mu, \theta)) \longrightarrow (\mathbb{T}(\mu'), L(\mu', \theta')).$$

Note that this notion of equivalence is formally in agreement with that used for pairs of tori and finite subgroups of a fixed order N .

In what follows, for any $A \in \mathrm{PGL}_2(\mathbb{Z})$, denote by A^{-T} the contragredient class i.e. the transformation defined by the inverse of the transpose of a matrix in the projective class of A : note that $(AB)^{-T} = A^{-T}B^{-T}$.

Proposition 3. *$\mathcal{F}(\mu, \theta)$ is isomorphic to $\mathcal{F}(\mu', \theta') \Leftrightarrow$ there exists $A \in \mathrm{PGL}_2(\mathbb{Z})$ such that*

$$\mu' = A(\mu) \quad \text{and} \quad \theta' = A^{-T}(\theta).$$

Proof. Assume first that $\mu, \mu' \in \mathbb{H}$ and $\mathcal{F}(\mu, \theta)$ and $\mathcal{F}(\mu', \theta')$ are isomorphic via the homothety defined by $\lambda \in \mathbb{C}$ with $\lambda \cdot \Lambda(\mu) = \Lambda(\mu')$. Then we have $\lambda\mu = a\mu' + b$, $\lambda = c\mu' + d$ where

$$B = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{Z}).$$

Thus $\mu = B(\mu')$ or $\mu' = B^{-1}(\mu)$. On the other hand,

$$\begin{aligned} \lambda \cdot \tilde{L}(\mu, \theta) &= \mathbb{R} \cdot ((c\mu' + d) + \theta(a\mu' + b)) \\ &= \mathbb{R} \cdot ((b\theta + d) + (a\theta + c)\mu') \\ &= \mathbb{R} \cdot (1 + B^T(\theta)\mu') \\ &= \tilde{L}(B^{-1}(\mu), B^T(\theta)). \end{aligned}$$

This shows that multiplication by λ induces an equivalence of foliations

$$\mathcal{F}(\mu, \theta) \longrightarrow \mathcal{F}(B^{-1}(\mu), B^T(\theta)).$$

Writing $A = B^{-1}$ we obtain the form of equivalence stated in the Proposition.

In case $(\mu', \theta') = (-\mu, -\theta)$ then we take

$$A = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z})$$

and noting that $A = A^T = A^{-1}$, we have $(A(\mu), A^{-T}(\theta)) = (-\mu, -\theta)$. The argument above is symmetric, so that if there exists $A \in \mathrm{PGL}_2(\mathbb{Z})$ such that $\mu' = A(\mu)$ and $\theta' = A^{-T}(\theta)$, then the corresponding Kronecker foliations are equivalent. \square

The moduli space of isomorphism classes of quantum tori is defined [27]

$$\mathrm{Mod}^{\mathrm{qt}} = \mathrm{PGL}_2(\mathbb{Z}) \backslash (\mathbb{R} - \mathbb{Q})$$

which may be viewed as a kind of boundary of the classical moduli space $\mathrm{Mod}^{\mathrm{cl}}$. Note that $\mathrm{Mod}^{\mathrm{qt}}$ is itself a non Hausdorff space since $\mathrm{PGL}_2(\mathbb{Z})$ acts densely on $\mathbb{R} - \mathbb{Q}$. The orbit of $[\infty]$ of \mathbb{Q} is viewed as the ideal boundary of $\mathrm{Mod}^{\mathrm{qt}}$ and we write as well

$$\overline{\mathrm{Mod}}^{\mathrm{qt}} := \mathrm{Mod}^{\mathrm{qt}} \cup [\infty].$$

By Proposition 2, the moduli space of generalized Kronecker foliations is the «signed» Anosov foliation

$$\mathrm{Mod}^{\mathrm{kf}} = \mathrm{PGL}_2(\mathbb{Z}) \backslash (\pm\mathbb{H} \times \mathbb{S}^1)$$

where $A \in \mathrm{PGL}_2(\mathbb{Z})$ acts by

$$(21) \quad A \cdot (\mu, \theta) = (A(\mu), A^{-T}(\theta)).$$

We regard the images of $\pm\mathbb{H} \times \{\theta\}$ in $\mathrm{Mod}^{\mathrm{kf}}$ as the leaves. This foliation fibers over $\mathrm{Mod}^{\mathrm{cl}} = \mathrm{PGL}_2(\mathbb{Z}) \backslash \pm\mathbb{H}$, in which the fiber $\mathrm{Mod}_{[i]}^{\mathrm{kf}}$ over the class $[i] \in \mathrm{Mod}^{\mathrm{cl}}$ parametrizes the «classical» Kronecker foliations. We have the following moduli space analogue of Proposition 2:

Proposition 4. *The leaf space of $\mathrm{Mod}^{\mathrm{kf}}$ is in canonical bijection with $\overline{\mathrm{Mod}}^{\mathrm{qt}}$.*

Proof. Since the fiber $\mathrm{Mod}_{[i]}^{\mathrm{kf}}$ is a complete transversal of the foliation $\mathrm{Mod}^{\mathrm{kf}}$, the leaf space of $\mathrm{Mod}^{\mathrm{kf}}$ may be identified with the set of leaf classes of elements of $\mathrm{Mod}_{[i]}^{\mathrm{kf}}$. The latter is the image of $\{i\} \times \mathbb{S}^1 \subset \pm\mathbb{H} \times \mathbb{S}^1$ under the suspension quotient. In particular, two points of $\mathrm{Mod}_{[i]}^{\mathrm{kf}}$ lie on the same leaf if and only if their preimages (i, θ) , $(i, \theta') \in \{i\} \times \mathbb{S}^1$ satisfy $\theta' = A(\theta)$ for some $A \in \mathrm{PGL}_2(\mathbb{Z})$ acting projective linearly on $\mathbb{S}^1 \approx \mathbb{R} \cup \{\infty\}$. Thus the leaf space of $\mathrm{Mod}^{\mathrm{kf}}$ may be put in canonical bijection with $\mathrm{PGL}_2(\mathbb{Z}) \backslash \mathbb{S}^1 \approx \overline{\mathrm{Mod}}^{\mathrm{qt}}$. \square

As mentioned in the Introduction, $\mathrm{Mod}^{\mathrm{kf}}$ provides a natural generalization of the moduli space $\Gamma_0(N) \backslash \mathbb{H}$ that classifies isomorphism classes of ordered pairs (E, C) , where E is an elliptic curve defined over \mathbb{C} , and C is a cyclic subgroup of E of order N .

We could extend j^{qt} – which is defined on the transversal $\mathrm{Mod}_{[i]}^{\mathrm{kf}}$ – to all of $\mathrm{Mod}^{\mathrm{kf}}$ using a similar definition to that found in §1, but the discontinuity and multivaluedness would persist. Instead we will use nonstandard models to construct an analogous space which fibers over $\mathrm{Mod}^{\mathrm{kf}}$ on which j^{qt} lifts to a continuous single valued function.

5. NONSTANDARD STRUCTURES

In what follows, \mathcal{S} is a discrete, infinite set.

Ultrafilters and Stone Spaces

Recall that a filter on \mathcal{S} is a subset $\mathfrak{f} \subset 2^{\mathcal{S}}$ not containing the empty set, which is closed with respect to finite intersections and upward inclusions ($X \in \mathfrak{f}$ and $Y \supset X \Rightarrow Y \in \mathfrak{f}$). Dually the set of complements $I_{\mathfrak{f}} := \{X \mid \mathcal{S} - X \in \mathfrak{f}\}$ is a proper ideal in the Boolean algebra $2^{\mathcal{S}}$. A maximal filter \mathfrak{u} is called an ultrafilter, whose set of complements $I_{\mathfrak{u}}$ is a maximal ideal of $2^{\mathcal{S}}$. A filter \mathfrak{f} is called nonprincipal if there exists no $X \in \mathfrak{f}$ with $X \subset Y$ for all $Y \in \mathfrak{f}$, or dually, if $I_{\mathfrak{f}}$ is a nonprincipal ideal. See [21].

If one has a family $\mathcal{A} \subset 2^{\mathcal{S}}$ of subsets not containing the empty set and satisfying the finite intersection property, there is a unique minimal filter containing \mathcal{A} , the filter $\langle \mathcal{A} \rangle$ generated by \mathcal{A} . For example, if \mathcal{S} is directed, $\gamma \in \mathcal{S}$ and $\hat{\gamma} = \{\gamma' \geq \gamma\}$ is the cone over γ , then by directedness $\mathcal{A} = \{\hat{\gamma}\}$ satisfies the finite intersection property and we will call

$$\mathfrak{c} = \mathfrak{c}_{\mathcal{S}} = \langle \mathcal{A} \rangle$$

the **cone filter** on \mathcal{S} . Note that \mathfrak{c} is nonprincipal: indeed, if there were a set X contained in all members of \mathfrak{c} , then for any $\gamma_0 \in X$ and $\gamma > \gamma_0$ we would have $X \not\subset \hat{\gamma} \in \mathfrak{c}$. An ultrafilter $\mathfrak{u} \supset \mathfrak{c}$ will be called a **cone ultrafilter**¹. Cone ultrafilters are nonprincipal. Moreover, every element $X \in \mathfrak{u}$ of a cone ultrafilter is a directed set, and so in particular, can be used to index nets.

The set of ultrafilters $\text{Ult}(\mathcal{S})$ on \mathcal{S} , equipped with the topology generated by the opens

$$V_X = \{\mathfrak{u} \mid X \in \mathfrak{u}\}, \quad X \in 2^{\mathcal{S}}$$

is called the Stone space of \mathcal{S} [22]. One has that $V_X^{\complement} = V_{X^{\complement}}$ where \complement means complement, so that the V_X are also closed. With this topology, $\text{Ult}(\mathcal{S})$ is totally-disconnected and compact, homeomorphic to the Stone-Cech compactification of \mathcal{S} or dually, to the space of maximal ideals $\text{Spec}(2^{\mathcal{S}})$ equipped with the dual Stone topology. The isolated points are the principal ultrafilters.

When \mathcal{S} is directed, the subspace $\text{Cone}(\mathcal{S})$ of cone ultrafilters is closed since

$$\text{Cone}(\mathcal{S}) = \bigcap_{\gamma \in \mathcal{S}} V_{\hat{\gamma}}.$$

In addition, $\text{Cone}(\mathcal{S})$ is perfect as all of its elements are nonprincipal ultrafilters, hence are non-isolated points. In particular, $\text{Cone}(\mathcal{S})$ is a (generalized) Cantor set, of cardinality possibly greater than that of the continuum.

Ultraproducts

Let L be a first order language, \mathcal{S} a directed set and $\{M_{\iota}\}_{\iota \in \mathcal{S}}$, a family of L -structures (e.g. a family of groups, rings, fields, etc) [19], [28]. Then the reduced product [8] of the M_{ι} with respect to \mathfrak{f} a filter on \mathcal{S} is the L -structure

$$[M_{\iota}]_{\mathfrak{f}} := \prod M_{\iota} / \sim_{\mathfrak{f}}$$

where $(x_{\iota}) \sim_{\mathfrak{f}} (x'_{\iota})$ if and only if $\{\iota \mid x_{\iota} = x'_{\iota}\} \in \mathfrak{f}$. If $M_{\iota} = M$ for all ι , the reduced product is denoted

$${}^*M_{\mathfrak{f}}$$

and called the reduced power of M with respect to \mathfrak{f} . If $\mathfrak{f} = \mathfrak{u}$ is an ultrafilter, the reduced product (reduced power) is called an ultraproduct (ultrapower).

¹The ultraproduct proof of the compactness theorem of first order logic uses a cone ultrafilter on $\mathcal{S} = \text{Fin}(T)$ where T is a finitely satisfiable first order theory [31].

By Łoś' Theorem [19], the ultrapower *M_u is an elementary extension of M , where the embedding $M \hookrightarrow {}^*M_u$ is given by the constant nets. What this means is that *M_u is a nonstandard model of M i.e. it satisfies the same set of first order L -sentences as M . In particular if M is a group, ring or field than so is *M_u . As one varies the ultrafilter, one obtains a sheaf

$${}^*\check{M} \rightarrow \text{Ult}(\mathcal{I})$$

whose fiber over u is *M_u , c.f. [24].

Note 5. If one assumes the Continuum Hypothesis (CH) and M is countable, then any two nonprincipal ultrafilters produce isomorphic ultrapowers [3]. More generally, if the complete theory of M is uncountably categorical – which is the case for $M = \mathbb{C}$ – then again assuming the CH, the nonprincipal fibers of ${}^*\check{M}$ will all be isomorphic, though not canonically so [28]. We will not, however, assume CH in this article.

If $\mathcal{I} = \mathbb{N}$ and u is a fixed nonprincipal ultrafilter on \mathbb{N} , then we will suppress the ultrafilter in our notation and denote the ultrapower

$${}^*M := {}^*M_u,$$

informally referring to it as «nonstandard M »; its elements will then be denoted *x , representatives of which are sequences $\{x_i\}$ in M .

Extended reals

We now turn to some specific ultrapowers which will be of interest to us: the nonstandard versions of the integers, the rationals, the reals and the complexes related in the usual way: ${}^*\mathbb{Z} \subset {}^*\mathbb{Q} \subset {}^*\mathbb{R} \subset {}^*\mathbb{C}$. Note that each of these structures contains classes corresponding to unbounded sequences, and are therefore non-Archimedean (as rings or as fields). In addition, ${}^*\mathbb{Z}$, ${}^*\mathbb{Q}$ and ${}^*\mathbb{R}$ are linearly ordered and the least upper bound property does not hold in ${}^*\mathbb{R}$ [32], [16].

It can be easily checked that the field ${}^*\mathbb{Q}$ is the field of fractions of the subring ${}^*\mathbb{Z}$. In addition, ${}^*\mathbb{Q}$ is also the field of fractions of another, *local* subring, defined as follows. Let $|\cdot|$ be the Archimedean absolute value on \mathbb{Q} . Then $|\cdot|$ induces in ${}^*\mathbb{Q}$ a nonstandard absolute value with values in ${}^*\mathbb{R}_+ =$ the nonnegative elements of ${}^*\mathbb{R}$. The set of bounded elements

$${}^*\mathbb{Q}_{\text{fin}} = \{ {}^*q \in {}^*\mathbb{Q} \mid \text{there exists } r \in \mathbb{R}_+ \text{ such that } |{}^*q| < r \}$$

is a local ring with maximal ideal the set of infinitesimals

$${}^*\mathbb{Q}_\varepsilon = \{ {}^*q \in {}^*\mathbb{Q} \mid \text{for all non-0 } r \in \mathbb{R}_+, |{}^*q| < r \}.$$

We shall write ${}^*x \simeq {}^*y$ whenever ${}^*x - {}^*y \in {}^*\mathbb{Q}_\varepsilon$ and say that *x and *y are asymptotic or infinitesimal to one another. We shall also refer to such a relation as an infinitesimal equation.

There is a canonical epimorphism

$$\text{std} : {}^*\mathbb{Q}_{\text{fin}} \longrightarrow \mathbb{R}$$

called the **standard part map**: for any ${}^*q \in {}^*\mathbb{Q}_{\text{fin}}$, $\text{std}({}^*q)$ is defined to be the unique accumulation point of any representative sequence $\{q_\alpha\}$ recognized by the ultrafilter. More precisely, for any representative sequence $\{q_\alpha\}$, there exists $X \in u$ such that $\{q_\alpha\}_X$ converges to a point $\text{std}({}^*q) \in \mathbb{R}$, which depends neither on $\{q_\alpha\}$ nor on X [32].

The kernel of std is ${}^*\mathbb{Q}_\varepsilon$ so that we have an isomorphism of fields

$${}^*\mathbb{Q}_{\text{fin}}/{}^*\mathbb{Q}_\varepsilon \cong \mathbb{R}.$$

One may compare this situation with that of the p -adic numbers \mathbb{Q}_p , where the quotient of the ring of integers by its maximal ideal is the finite field \mathbb{F}_p with p elements.

Extending $|\cdot|$ to ${}^*\mathbb{R}$, we define in the same way the local ring ${}^*\mathbb{R}_{\text{fin}}$ with maximal ideal ${}^*\mathbb{R}_\varepsilon$ obtaining ${}^*\mathbb{R}_{\text{fin}}/{}^*\mathbb{R}_\varepsilon \cong \mathbb{R}$. We may similarly recover \mathbb{C} from the quotient ${}^*\mathbb{C}_{\text{fin}}/{}^*\mathbb{C}_\varepsilon$ where ${}^*\mathbb{C}_{\text{fin}}, {}^*\mathbb{C}_\varepsilon$ are defined using the usual absolute value in \mathbb{C} .

The quotient

$${}^*\mathbb{R} := {}^*\mathbb{R}/{}^*\mathbb{R}_\varepsilon \cong {}^*\mathbb{Q}/{}^*\mathbb{Q}_\varepsilon$$

is a real vector space (but not a topological vector space with respect to the quotient order topology) which we shall call the **extended reals** [11], [12]. Note that ${}^*\mathbb{R}$ contains \mathbb{R} canonically, and also ${}^*\mathbb{Z}$ since ${}^*\mathbb{Z} \cap {}^*\mathbb{R}_\varepsilon = \{0\}$. We will view ${}^*\mathbb{R}$ as «foliated» by the cosets ${}^*x + \mathbb{R}$. The subring ${}^*\mathbb{Z}$ defines a transversal (in the sense that it has non trivial and discrete intersection with each coset leaf ${}^*x + \mathbb{R}$) and the «leaf space» may be identified with ${}^*\mathbb{Z}/\mathbb{Z}$, which a priori is not endowed with any particular topology. We define the extended complex numbers ${}^*\mathbb{C}$ in exactly the same way.

6. DIOPHANTINE APPROXIMATION GROUPS

Fix ${}^*\mathbb{Z}$ a nonstandard ring of integers and let $(\mu, \theta) \in \pm\mathbb{H} \times \mathbb{S}^1$. Since $\Lambda(\mu) \subset \mathbb{C}$ is discrete, the ultrapower ${}^*\Lambda(\mu)$ is naturally a subgroup of the vector space ${}^*\mathbb{C} = {}^*\mathbb{C}/{}^*\mathbb{C}_\varepsilon$. In the Proposition which follows, we endow ${}^*\mathbb{C}$ with the euclidean topology along its coset leaves ${}^*z + \mathbb{C}$ and the discrete topology transversally.

Proposition 5. *The quotient*

$${}^*\mathbb{C}/{}^*\Lambda(\mu)$$

is a topological group topologically isomorphic to $\mathbb{T}(\mu)$.

Proof. Note that $({}^*\mathbb{C}, +)$ is a topological group. In addition, ${}^*\Lambda(\mu)$ is a complete transversal for ${}^*\mathbb{C}$, so that every ${}^*z \in {}^*\mathbb{C}$ can be translated by an element of ${}^*\Lambda(\mu)$ to $\mathbb{C} \subset {}^*\mathbb{C}$. Since ${}^*\Lambda(\mu) \cap \mathbb{C} = \Lambda(\mu)$, then ${}^*\mathbb{C}/{}^*\Lambda(\mu) = \mathbb{C}/\Lambda(\mu)$ and the Proposition follows. \square

Define the **extended line of μ -slope θ** as

$${}^*\tilde{L}(\mu, \theta) := {}^*\mathbb{R} \cdot (\theta\mu + 1) \subset {}^*\mathbb{C}.$$

Definition 1. We say that ${}^*n \in {}^*\mathbb{Z}$ is a **diophantine approximation** of θ (relative to μ) if there exists ${}^*m \in {}^*\mathbb{Z}$ such that

$$(22) \quad {}^*m\mu + {}^*n \in {}^*\tilde{L}(\mu, \theta) \cap {}^*\Lambda(\mu).$$

The next Proposition shows that the condition (22) depends only on θ .

Proposition 6. *The pair $({}^*m, {}^*n)$ defines a diophantine approximation of θ relative to $\mu \Leftrightarrow$ its coordinates satisfy (in ${}^*\mathbb{R}$)*

$$(23) \quad {}^*n\theta = {}^*m$$

Proof. The argument involves simple manipulations of equations in ${}^*\mathbb{R}$ using its \mathbb{R} -vector space structure. First, *n is a diophantine approximation of θ relative to $\mu = a + ib \Leftrightarrow$ there exists ${}^*r \in {}^*\mathbb{R}$ such that ${}^*m\mu + {}^*n = {}^*r(\theta\mu + 1)$. Separating into real and imaginary parts gives

$$(24) \quad {}^*ma + {}^*n = {}^*r(\theta a + 1) \quad \text{and} \quad {}^*mb = {}^*r\theta b.$$

The second equation of (24) in turn yields ${}^*m = {}^*r\theta$, which, when plugged back into the first equation of (24), gives ${}^*r\theta a + {}^*n = {}^*r\theta a + {}^*r$ or ${}^*r = {}^*n$. Plugging the latter into ${}^*m = {}^*r\theta$ gives (23). Conversely, if $({}^*m, {}^*n)$ satisfies (23), then taking ${}^*r = {}^*n$ gives the pair of equations (24), which imply the condition (22). \square

The absolute version of diophantine approximation (23) is that used in [11], [13]. It is clear from the form of (23) that the collection of diophantine approximations of θ relative to μ forms a subgroup of ${}^*\mathbb{Z}$ denoted

$${}^*\mathbb{Z}(\theta),$$

which is independent of μ . Note that this group is uncountably infinite and torsion-free.

Theorem 5. *The group ${}^*\mathbb{Z}(\theta)$ is an ideal in ${}^*\mathbb{Z} \Leftrightarrow \theta \in \mathbb{Q}$. If $\theta, \theta' \in \mathbb{S}^1$ satisfy $A(\theta) = \theta'$ for some $A \in \text{PGL}_2(\mathbb{Z})$ then ${}^*\mathbb{Z}(\theta) \cong {}^*\mathbb{Z}(\theta')$.*

This is proved in [11]; for the convenience of the reader, we include here a

Proof. If $\theta \in \mathbb{Q}$, then a pair ${}^*n, {}^*m$ satisfies (23) \Leftrightarrow we have the equality in the field ${}^*\mathbb{R}$: ${}^*n\theta = {}^*m$. Such an equality is invariant with respect to multiplication by elements of ${}^*\mathbb{Z}$, which shows that ${}^*\mathbb{Z}(\theta)$ is an ideal. If $\theta \in \mathbb{R} - \mathbb{Q}$ and ${}^*n \in {}^*\mathbb{Z}(\theta)$ then by irrationality we can find ${}^*N \in {}^*\mathbb{Z}$ such that ${}^*N{}^*n\theta$ contains a representative sequence asymptotic mod \mathbb{Z} to any element of \mathbb{S}^1 we choose. If this element is not 0, then ${}^*N{}^*n \notin {}^*\mathbb{Z}(\theta)$, showing that ${}^*\mathbb{Z}(\theta)$ is not an ideal. Now let

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

be a (representative of an) element of $\text{PGL}_2(\mathbb{Z})$. If $({}^*m, {}^*n)$ satisfies (23) then

$$\begin{pmatrix} {}^*m' \\ {}^*n' \end{pmatrix} = A \begin{pmatrix} {}^*m \\ {}^*n \end{pmatrix} = \begin{pmatrix} a{}^*m + b{}^*n \\ c{}^*m + d{}^*n \end{pmatrix}$$

satisfies the analogue of (23) for $A(\theta)$. Indeed, we have in ${}^*\mathbb{R}$ that ${}^*n'A(\theta) = {}^*m' \Leftrightarrow {}^*n'(a\theta + b) = {}^*m'(c\theta + d) \Leftrightarrow (c{}^*m + d{}^*n)(a\theta + b) = (a{}^*m + b{}^*n)(c\theta + d)$. But the latter equation is equivalent in ${}^*\mathbb{R}$ to ${}^*n\theta = {}^*m$. \square

The element *m associated to *n is unique: we refer to it as the dual of *n and use the notation

$${}^*n^\perp := {}^*m.$$

The set of duals ${}^*\mathbb{Z}^\perp(\theta)$ is a group, and when $\theta \neq 0$, it is canonically isomorphic to ${}^*\mathbb{Z}(\theta)$ and equal to ${}^*\mathbb{Z}(\theta^{-1})$. In addition,

$${}^*\Lambda(\mu, \theta) = \left\{ \mu \cdot {}^*n^\perp + {}^*n \mid {}^*n \in {}^*\mathbb{Z}(\theta) \right\}$$

defines a subgroup of ${}^*\Lambda(\mu)$ called the **group of (μ, θ) -fractions**, and the map ${}^*n \mapsto \mu \cdot {}^*n^\perp + {}^*n$ defines an isomorphism ${}^*\mathbb{Z}(\theta) \cong {}^*\Lambda(\mu, \theta)$.

The following Proposition expresses ${}^*\Lambda(\mu, \theta)$ as the intersection of a pair of ultrapowers of standard lattices of rank 2 resp. 1. Given $\nu \in \pm\mathbb{H}$, let

$${}^*\Lambda_{\Delta}(\nu) = \{ {}^*n(1+\nu) \mid n \in {}^*\mathbb{Z} \} \subset {}^*\Lambda(\nu)$$

be the ultrapower of the group $\mathbb{Z} \cdot (1+\nu) \subset \Lambda(\nu)$ that uniformizes the diagonal cycle $c(\nu) \subset \mathbb{T}(\nu)$.

Proposition 7. *For any $\theta \in \mathbb{R}$, $\mu \in \pm\mathbb{H}$,*

$${}^*\Lambda(\mu, \theta) = {}^*\Lambda(\mu) \cap {}^*\Lambda_{\Delta}(\theta\mu).$$

Proof. By Proposition 6, if ${}^*m\mu + {}^*n \in {}^*\Lambda(\mu, \theta)$ then ${}^*m\mu + {}^*n = {}^*n(\mu\theta + 1) \in {}^*\Lambda_{\Delta}(\theta\mu)$. Conversely if ${}^*m\mu + {}^*n = {}^*n'(\mu\theta + 1) \in {}^*\Lambda(\mu) \cap {}^*\Lambda_{\Delta}(\theta\mu)$ then ${}^*m\mu + {}^*n \in {}^*\tilde{L}(\mu, \theta) \cap {}^*\Lambda(\mu)$. \square

There is a natural homomorphism of abelian groups

$$\text{std}(\mu, \theta) : {}^*\tilde{L}(\mu, \theta) \longrightarrow \mathbb{T}(\mu)$$

defined as follows. Take a representative sequence $\{r_{\alpha}\} \in {}^*r \in {}^*\mathbb{R}$, and consider the image in $\mathbb{T}(\mu)$ of the sequence

$$\{r_{\alpha} \cdot (\theta\mu + 1)\}$$

in the leaf $L(\mu, \theta) \subset \mathbb{T}(\mu)$ through the origin. Since $\mathbb{T}(\mu)$ is compact, the ultrafilter will recognize a unique limit point of this sequence, which is independent of the choice of $\{r_{\alpha}\} \in {}^*r$ (again, see [32] for more on this compactness principle). We define

$$\text{std}(\mu, \theta)(({}^*r \cdot (\theta\mu + 1)))$$

to be this limit point.

Notice that the leaves of ${}^*\mathbb{R}$ gives rise to leaves of ${}^*\tilde{L}(\mu, \theta)$, defined as the scalar multiples $(\theta\mu + 1) \cdot ({}^*r + \mathbb{R})$. We note that the leaf corresponding to ${}^*r = 0$ is the line $\tilde{L}(\mu, \theta) \subset \mathbb{C}$ which was defined in §2: that is we have $\tilde{L}(\mu, \theta) \subset {}^*\tilde{L}(\mu, \theta)$. The map $\text{std}(\mu, \theta)$ transports these leaves to the leaves of the associated Kronecker foliation $\mathcal{F}(\mu, \theta)$.

Theorem 6. *If $\theta \in \mathbb{R} - \mathbb{Q}$ then $\text{std}(\mu, \theta)$ is surjective with kernel ${}^*\Lambda(\mu, \theta)$.*

Proof. Surjectivity follows from the density of $L(\mu, \theta)$ in $\mathbb{T}(\mu)$. The map $\text{std}(\mu, \theta)$ coincides with the restriction of the epimorphism ${}^*\mathbb{C} \rightarrow \mathbb{T}(\mu)$ of Proposition 5 to the subspace ${}^*\tilde{L}(\mu, \theta)$, so that the kernel is ${}^*\tilde{L}(\mu, \theta) \cap {}^*\Lambda(\mu) = {}^*\Lambda(\mu, \theta)$. \square

Thus we have the «foliated group» isomorphisms

$${}^*\mathbb{R}/{}^*\mathbb{Z}(\theta) \cong {}^*\tilde{L}(\mu, \theta)/{}^*\Lambda(\mu, \theta) \cong \mathcal{F}(\mu, \theta).$$

Here we point out that the last isomorphism is not topological: nevertheless it is possible to put on ${}^*\tilde{L}(\mu, \theta)$ a new transverse topology so that the action of ${}^*\Lambda(\mu, \theta)$ is by homeomorphisms, and that the quotient ${}^*\tilde{L}(\mu, \theta)/{}^*\Lambda(\mu, \theta)$ becomes a foliation isomorphic to $\mathcal{F}(\mu, \theta)$, see [11], [13]. By Proposition 7, there are «covering maps»

$$\mathcal{F}(\mu, \theta) \rightarrow \mathbb{T}(\mu) \quad \text{and} \quad \mathcal{F}(\mu, \theta) \rightarrow c(\theta\mu)$$

where $c(\theta\mu) \subset \mathbb{T}(\theta\mu)$ is the diagonal cycle.

7. ULTRASOLENOIDS

As in previous sections we denote by ${}^*\mathbb{Z}$, ${}^*\mathbb{C}$ the ultrapowers of \mathbb{Z} , \mathbb{C} with respect to a fixed nonprincipal ultrafilter on \mathbb{N} . Let S be a set, *S_u be an arbitrary ultrapower with respect to some index set \mathcal{A} and ultrafilter u . Recall that a hyperfinite subset [16], [32] of *S_u is an ultraproduct of the form

$$[F_\alpha]_u \subset [S_\alpha = S]_u = {}^*S_u$$

where $F_\alpha \subset S$ is finite for all $\alpha \in \mathcal{A}$. Thus elements of $[F_\alpha]$ are classes of sequences $\{x_\alpha\}$ for which $x_\alpha \in F_\alpha$ for all α . Note that every finite subset of *S_u is hyperfinite.

Consider now the directed set (directed by inclusion)

$$\mathcal{H} = \{\text{hyperfinite subsets } [F_\alpha] \subset {}^*\mathbb{Z}^2 - \{0,0\}\}.$$

Denote by \mathfrak{c} the cone filter on \mathcal{H} , which we recall was defined in §3 as the (nonprincipal) filter generated by the cones

$$\text{cone}([F_\alpha]) = \{[F'_\alpha] \mid [F_\alpha] \subset [F'_\alpha]\}.$$

Let $\text{Ult}(\mathcal{H})$ be the Stone space of ultrafilters on \mathcal{H} , and denote by $\text{Cone}(\mathcal{H}) \subset \text{Ult}(\mathcal{H})$ the subspace of ultrafilters extending \mathfrak{c} . Each element $u \in \text{Cone}(\mathcal{H})$ is nonprincipal, so $\text{Cone}(\mathcal{H})$ is a Cantor set. For us the importance of the cone ultrafilters is that they will provide partial summation schemes that correspond well to the classical definition of a convergent infinite series.

Following [24], we define a sheaf ${}^\circ\check{\mathbb{C}}$ over $\text{Ult}(\mathcal{H})$ as follows: for each $u \in \text{Ult}(\mathcal{H})$, the stalk over u , ${}^\circ\mathbb{C}_u$, is the ultrapower of ${}^*\mathbb{C}$ with respect to u . Let ${}^\circ\check{\Gamma}$ be the ${}^*\mathbb{C}$ -algebra of set-theoretic sections of ${}^\circ\check{\mathbb{C}}$: the ${}^*\mathbb{C}$ -algebra structure comes from the fact that ${}^*\mathbb{C}$ is canonically included in each fiber ${}^\circ\mathbb{C}_u$ via the constant net inclusion. In particular, we have canonical \mathbb{C} -algebra inclusions $\mathbb{C} \subset {}^*\mathbb{C} \subset {}^\circ\check{\Gamma}$ defined by the constant sections. Finally, denote by ${}^\circ\check{\mathbb{C}}^{\text{cone}}$ the restriction of ${}^\circ\check{\mathbb{C}}$ to $\text{Cone}(\mathcal{H})$ and by ${}^\circ\check{\Gamma}^{\text{cone}}$ the sections of ${}^\circ\check{\mathbb{C}}^{\text{cone}}$. There is a canonical ${}^*\mathbb{C}$ -algebra epimorphism ${}^\circ\check{\Gamma} \rightarrow {}^\circ\check{\Gamma}^{\text{cone}}$ given by restriction.

We now define subsheaves that correspond to $\theta \in \bar{\mathbb{R}}$. For $\theta \neq \infty$ let

$${}^*\mathbb{Z}^2(\theta) = \left\{ ({}^*n^\perp, {}^*n) \mid {}^*n \in {}^*\mathbb{Z}(\theta) \right\} \subset {}^*\mathbb{Z}^2.$$

For $\theta = \infty$ we define ${}^*\mathbb{Z}^2(\infty) := {}^*\mathbb{Z} \times \{0\}$. Let $\mathcal{H}(\theta) \subset \mathcal{H}$ be the subset of hyperfinite subsets contained in ${}^*\mathbb{Z}^2(\theta)$. Let $\mathfrak{c}(\theta)$ be the cone filter of $\mathcal{H}(\theta)$. Denote by

$$\text{Cone}(\mathcal{H})(\theta) \subset \text{Ult}(\mathcal{H})$$

the subspace of ultrafilters u of \mathcal{H} (not of $\mathcal{H}(\theta)$) that extend $\mathfrak{c}(\theta)$. The ultrafilters belonging to $\text{Cone}(\mathcal{H})(\theta)$ are those ultrafilters of \mathcal{H} that observe the group ${}^*\mathbb{Z}^2(\theta)$.

Let ${}^\circ\check{\mathbb{C}}^{\text{cone}}(\theta)$ be the restriction of ${}^\circ\check{\mathbb{C}}$ to $\text{Cone}(\mathcal{H})(\theta)$ and let ${}^\circ\check{\Gamma}^{\text{cone}}(\theta)$ be its ${}^*\mathbb{C}$ -algebra of sections. The restriction map gives an algebra epimorphism ${}^\circ\check{\Gamma} \rightarrow {}^\circ\check{\Gamma}^{\text{cone}}(\theta)$. The Lemma which follows shows that the sheaves ${}^\circ\check{\mathbb{C}}^{\text{cone}}$, ${}^\circ\check{\mathbb{C}}^{\text{cone}}(\theta)$ are disjoint for all $\theta \in \bar{\mathbb{R}}$.

Lemma 2. *Let $\theta, \eta \in \bar{\mathbb{R}}$ be distinct. Then*

$$\text{Cone}(\mathcal{H})(\theta) \cap \text{Cone}(\mathcal{H})(\eta) = \emptyset = \text{Cone}(\mathcal{H})(\theta) \cap \text{Cone}(\mathcal{H})(\eta).$$

Proof. Suppose that $u \in \text{Cone}(\mathcal{H})(\theta) \cap \text{Cone}(\mathcal{H})(\eta)$ so that u contains both \mathfrak{c} and $\mathfrak{c}(\theta)$. Let $F \subset \mathbb{Z}^2 \subset {}^*\mathbb{Z}^2$ be a finite set containing a non zero element $(m, n) \notin {}^*\mathbb{Z}^2(\theta)$ (if $\theta \in \mathbb{R} - \mathbb{Q}$ this is true of any non-zero (m, n)). Then $\text{cone}(F) \in \mathfrak{c} \subset u$ and each element of $\text{cone}(F)$

is a subset of ${}^*\mathbb{Z}^2$ which contains (m, n) . On the other hand, for any $X \in \mathfrak{c}(\theta) \subset \mathfrak{u}$, X cannot contain any hyper finite subsets which contain (m, n) . In particular we must have that $X \cap \text{cone}(F) = \emptyset$ is empty, contradicting the fact that \mathfrak{u} is an ultrafilter. Now for all $\theta \neq \eta$ we have ${}^*\mathbb{Z}^2(\theta) \cap {}^*\mathbb{Z}^2(\eta) = (0, 0)$. Indeed, if $({}^*n^\perp, {}^*n) \in {}^*\mathbb{Z}^2(\theta)$ then $\theta \simeq {}^*n^\perp/{}^*n$ so that it is not possible to also have $\eta \simeq {}^*n^\perp/{}^*n$ for $\theta \neq \eta$. In particular $\mathcal{H}(\theta) \cap \mathcal{H}(\eta) = \{(0, 0)\}$ and therefore $\text{Cone}(\mathcal{H})(\theta) \cap \text{Cone}(\mathcal{H})(\eta) = \emptyset$. \square

We define actions of $\text{GL}_2(\mathbb{Z})$ on the sheaves just considered, as well as on their algebras of sections. First note that the left action of $\text{GL}_2(\mathbb{Z})$ on ${}^*\mathbb{Z}^2 - \{0, 0\}$ induces one on hyperfinite sets, $[F_\alpha] \mapsto [AF_\alpha]$ for $A \in \text{GL}_2(\mathbb{Z})$. This in turn induces an action on $\text{Ult}(\mathcal{H})$ which preserves $\text{Cone}(\mathcal{H})$ and identifies $\text{Cone}(\mathcal{H})(\theta)$ with $\text{Cone}(\mathcal{H})(A(\theta))$.

We can now define an action of $\text{GL}_2(\mathbb{Z})$ on the sheaf ${}^\diamond\check{\mathbb{C}}$ as follows: if $\{{}^*z_{[F_\alpha]}\}$ represents an element of ${}^\diamond z \in {}^\diamond\mathbb{C}_{\mathfrak{u}}$ then

$$\{{}^*w_{[F_\alpha]}\} := \{{}^*z_{[AF_\alpha]}\}$$

represents an element of ${}^\diamond\mathbb{C}_{A^{-1}\mathfrak{u}}$. Indeed, suppose that $\{{}^*z'_{[F_\alpha]}\}$ is another net representing ${}^\diamond z$. Then there is a set $X \in \mathfrak{u}$ of hyperfinite sets such that $\{{}^*z_{[F_\alpha]}\}|_X = \{{}^*z'_{[F_\alpha]}\}|_X$. It follows that

$$\{{}^*w_{[F_\alpha]}\}|_{A^{-1}X} = \{{}^*z_{[F_\alpha]}\}|_X = \{{}^*z'_{[F_\alpha]}\}|_X = \{{}^*w'_{[F_\alpha]}\}|_{A^{-1}X}.$$

Therefore $\{{}^*w_{[F_\alpha]}\}$ and $\{{}^*w'_{[F_\alpha]}\}$ are equivalent modulo $A^{-1}\mathfrak{u}$.

Denote the action by $A \in \text{GL}_2(\mathbb{Z})$ defined in the previous paragraph by $A \circ {}^\diamond z$: we emphasize that it is *not* the matrix action along fibers. Rather, the action is a shift, and so acts by ${}^*\mathbb{C}$ -algebra isomorphisms along the fibers of ${}^\diamond\check{\mathbb{C}}$ fixing the constant net classes. That is, if ${}^\diamond z = {}^*z \in {}^\diamond\mathbb{C}_{\mathfrak{u}}$ is a constant net class then $A \circ {}^*z = {}^*z$ (viewed as an element of ${}^\diamond\mathbb{C}_{A^{-1}\mathfrak{u}}$). In particular, we obtain ${}^*\mathbb{C}$ -algebra isomorphisms $A : {}^\diamond\mathbb{C}_{A\mathfrak{u}} \rightarrow {}^\diamond\mathbb{C}_{\mathfrak{u}}$ for each $\mathfrak{u} \in \text{Ult}(\mathcal{H})$. This action stabilizes ${}^\diamond\check{\mathbb{C}}^{\text{cone}}$ and maps ${}^\diamond\check{\mathbb{C}}^{\text{cone}}(A(\theta))$ isomorphically onto ${}^\diamond\check{\mathbb{C}}^{\text{cone}}(\theta)$.

There is also an induced action by $A \in \text{GL}_2(\mathbb{Z})$ on elements of ${}^\diamond\check{\Gamma}$ defined

$$f \mapsto g = A \circ f, \quad g(\mathfrak{u}) := A \circ (f(A\mathfrak{u}))$$

which defines a ${}^*\mathbb{C}$ -isomorphism of ${}^\diamond\check{\Gamma}$ (since it acts as the identity on the constant sections ${}^*\mathbb{C} \subset {}^\diamond\check{\Gamma}$). Again, ${}^\diamond\check{\Gamma}^{\text{cone}}$ is preserved by this action and ${}^\diamond\check{\Gamma}^{\text{cone}}(A(\theta))$ is identified with ${}^\diamond\check{\Gamma}^{\text{cone}}(\theta)$.

We now form the quotient of ${}^\diamond\check{\mathbb{C}}$ with respect to the $\text{GL}_2(\mathbb{Z})$ action:

$${}^\diamond\hat{\mathbb{C}} := \text{GL}_2(\mathbb{Z}) \backslash {}^\diamond\check{\mathbb{C}}.$$

The result, as such, is no longer a sheaf but rather a solenoid-like object, in a sense made precise in *Note 6* below. We thus call ${}^\diamond\hat{\mathbb{C}}$ an **ultrasolenoid**. The quotient

$${}^\diamond\hat{\Gamma} := \text{GL}_2(\mathbb{Z}) \backslash {}^\diamond\check{\Gamma}$$

is called the algebra of **ultratransversals**: since $\text{GL}_2(\mathbb{Z})$ acts as the identity on the constant sections, ${}^\diamond\hat{\Gamma}$ acquires the structure of a ${}^*\mathbb{C}$ -algebra extension of ${}^*\mathbb{C}$.

In view of the fact that $\text{GL}_2(\mathbb{Z})$ stabilizes $\text{Cone}(\mathcal{H})$, we also have a subultrasolenoid

$$\hat{\mathbb{C}}^{\text{cl}} := \text{GL}_2(\mathbb{Z}) \backslash {}^\diamond\check{\mathbb{C}}^{\text{cone}} \subset {}^\diamond\hat{\mathbb{C}}$$

which we call the **classical ultrasolenoid** with the associated ultratransversal algebra:

$${}^\diamond\hat{\Gamma}^{\text{cl}} := \text{GL}_2(\mathbb{Z}) \backslash {}^\diamond\check{\Gamma}^{\text{cone}}.$$

The restriction map $\diamond\check{\Gamma} \rightarrow \diamond\check{\Gamma}^{\text{cone}}$ induces a projection

$$\pi^{\text{cl}} : \diamond\hat{\Gamma} \longrightarrow \diamond\hat{\Gamma}^{\text{cl}}.$$

Moreover, each equivalence class $[\theta] \in \text{GL}_2(\mathbb{Z}) \backslash \mathbb{R}$ gives rise as well to a subultra-solenoid

$$\hat{\mathbb{C}}^{\text{qt}}([\theta]) := \text{GL}_2(\mathbb{Z}) \backslash \left(\bigsqcup_{A \in \text{GL}_2(\mathbb{Z})} \diamond\check{\mathbb{C}}^{\text{cone}}(A\theta) \right) \subset \diamond\hat{\mathbb{C}}$$

which we call the **$[\theta]$ -quantum ultrasolenoid**, and a corresponding algebra of ultra-transversals

$$\hat{\Gamma}^{\text{qt}}([\theta]) := \text{GL}_2(\mathbb{Z}) \backslash \left(\bigsqcup_{A \in \text{GL}_2(\mathbb{Z})} \diamond\check{\Gamma}^{\text{cone}}(A\theta) \right).$$

The restriction map induces again a projection

$$\pi^{\text{qt}} : \diamond\hat{\Gamma} \longrightarrow \diamond\hat{\Gamma}^{\text{qt}}([\theta]).$$

The motive for forming these $\text{GL}_2(\mathbb{Z})$ quotients is to ensure that the «Eisenstein objects» we define in the sequel are automorphic.

Note 6. The ultrasolenoids defined above are sheaf theoretic generalizations of the classical solenoid $\hat{\mathbb{S}} = (\mathbb{R} \times \hat{\mathbb{Z}})/\mathbb{Z}$, where $\hat{\mathbb{Z}}$ is the profinite completion of \mathbb{Z} (a Cantor group) and the action is diagonal. We refer to the images of the fibers $\diamond\mathbb{C}_u$ as the «leaves» of $\diamond\hat{\mathbb{C}}$ (each of which is foliated by its $\ast\mathbb{C}$ -cosets). Recall from *Note 5* that assuming the Continuum Hypothesis, the leaves of $\diamond\hat{\mathbb{C}}$ will be isomorphic to one another, though not canonically so. This motivates the foliated view of these quotients. In particular, the elements of $\diamond\hat{\Gamma}$ are complete transversals of $\diamond\hat{\mathbb{C}}$.

We describe briefly the context in which the constructions given in this section will be used to define modular invariants. As described in the above paragraphs we have a pair of epimorphisms

$$\begin{array}{ccc} & \diamond\hat{\Gamma} & \\ \pi^{\text{cl}} \swarrow & & \searrow \pi^{\text{qt}}([\theta]) \\ \diamond\hat{\Gamma}^{\text{cl}} & & \diamond\hat{\Gamma}^{\text{qt}}([\theta]) \end{array}$$

We will define first in §9 the universal modular invariant as a (set-theoretic) function

$$\diamond\hat{j} : \pm\mathbb{H} \longrightarrow \diamond\hat{\Gamma}$$

for which

- the function $\diamond\hat{j}^{\text{cl}}(\mu) := \pi^{\text{cl}} \circ \diamond\hat{j}(\mu)$ yields a function asymptotic to the usual modular invariant of μ .
- the function $\diamond\hat{j}^{\text{qt}}(\mu, \theta) := \pi^{\text{qt}}([\theta]) \circ \diamond\hat{j}(\mu)$ defines the (nonstandard) quantum modular invariant of (μ, θ) . When $\mu = i$, the result will be asymptotic to a multimap containing the standard quantum modular invariant defined in §1; a slight modification gives j^{qt} exactly.

We will then reinterpret the universal modular invariant as a *continuous* function

$$\diamond j : \widehat{\diamond\text{Mod}} \longrightarrow \diamond\hat{\mathbb{C}},$$

where $\widehat{\diamond\text{Mod}}$ is a topological ultrasolenoid that we will define in the last part of §9.

8. EISENSTEIN ULTRATRANSVERSALS

We continue to fix as before ultrapowers ${}^*\mathbb{Z} \subset {}^*\mathbb{Q} \subset {}^*\mathbb{R} \subset {}^*\mathbb{C}$. In this section we associate to every $k \in \mathbb{Z}$ and each pair $(\mu, \theta) \in \pm\mathbb{H} \times \mathbb{S}^1$ an analogue of the classical Eisenstein series, defined as an ultratransversal

$$\diamond \hat{G}_k^{\text{qt}}(\mu, \theta) \in \diamond \hat{\Gamma}^{\text{qt}}.$$

Fix $\mu \in \pm\mathbb{H}$. For each hyperfinite set $[F_\alpha] \in \mathcal{H}$ and $k \in \mathbb{Z}$ consider the hyperfinite sum [16]

$$G_k(\mu)_{[F_\alpha]} = \sum_{[F_\alpha]} ({}^*m\mu + {}^*n)^{-2k} := \text{*class of } \left\{ \sum_{(m_\alpha, n_\alpha) \in F_\alpha} (m_\alpha \mu + n_\alpha)^{-2k} \right\} \in {}^*\mathbb{C}.$$

Note that this expression is well-defined even for $k < 2$, in contrast with to the classical situation. For example, when $k = 0$, we have

$$G_0(\mu)_{[F_\alpha]} = \text{hypercardinality of } [F_\alpha] = \text{*class of } \{|F_\alpha|\}.$$

The \mathcal{H} -net

$$(25) \quad \{G_k(\mu)_{[F_\alpha]}\}$$

defines as described in the previous section an element $\diamond \hat{G}_k(\mu) \in \diamond \check{\Gamma}$. We thus obtain a function

$$\diamond \hat{G}_k : \pm\mathbb{H} \longrightarrow \diamond \hat{\Gamma}.$$

Proposition 8. $\diamond \hat{G}_k$ is a modular form of weight k :

$$(A'(\mu))^k \cdot \diamond \hat{G}_k(A\mu) = \diamond \hat{G}_k(\mu)$$

for all $A \in \text{GL}_2(\mathbb{Z})$.

Proof. We will show that in $\diamond \check{\Gamma}$

$$(A'(\mu))^k \cdot \diamond \check{G}_k(A\mu) = A^T \circ \diamond \check{G}_k(\mu).$$

We calculate at the level of the net (25): for

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{Z})$$

we have

$$\begin{aligned} (A'(\mu))^k \cdot G_k(A\mu)_{[F_\alpha]} &= (c\mu + d)^{-2k} \sum_{[F_\alpha]} \left({}^*m \left(\frac{a\mu + b}{c\mu + d} \right) + {}^*n \right)^{-2k} \\ &= \sum_{[F_\alpha]} \left((a{}^*m + c{}^*n)\mu + (b{}^*m + d{}^*n) \right)^{-2k} \\ &= G_k(\mu)_{A^T[F_\alpha]} \end{aligned}$$

from which the statement follows. \square

Denote by

$$\diamond \hat{G}_k^{\text{cl}} : \pm\mathbb{H} \longrightarrow \diamond \hat{\Gamma}^{\text{cl}}$$

the composition of $\diamond \hat{G}_k$ with the projection $\pi^{\text{cl}} : \diamond \hat{\Gamma} \longrightarrow \diamond \hat{\Gamma}^{\text{cl}}$ defined in §7. For $k \geq 2$ and $\mu \in \pm\mathbb{H}$, let $G_k(\mu)$ be the usual (standard) Eisenstein series. Since $\mathbb{C} \subset \diamond \hat{\Gamma}^{\text{cl}}$ we may view G_k as defining a family of «constant ultratransversals»

$$G_k : \mathbb{H} \rightarrow \diamond \hat{\Gamma}^{\text{cl}}.$$

In what follows, for any pair of sections $f, g : \text{Ult}(\mathcal{H}) \rightarrow \diamond\check{\mathbb{C}}$ we write $f \simeq g$ if $f - g \in {}^*C_\varepsilon \subset {}^*C \subset \diamond C_u$ for all u . Notice that this relation is preserved by the action of $\text{GL}_2(\mathbb{Z})$, giving rise to the relation of infinitesimality of ultratransversals in $\diamond\hat{\Gamma}^{\text{cl}}$.

Proposition 9. *For all $\mu \in \pm\mathbb{H}$,*

$$G_k(\mu) \simeq \diamond\hat{G}_k(\mu)^{\text{cl}}.$$

Proof. Let $u \in \text{Cone}(\mathcal{H})$. It will be enough to check that for any finite subset $F \subset \mathbb{Z}^2$ that the net of hyperfinite sums over elements in $\text{cone}(F)$ converges to $G_k(\mu)$. This is certainly true if we restrict to the subnet of all *finite* subsets $F' \supset F$, because the classical Eisenstein series converges. Now if $[F_\alpha] \supset F$ is a general hyperfinite containing F , and $\varepsilon > 0$, let $F' \supset F$ be such that $G_k(\mu)_{F''}$ is ε -close to $G_k(\mu)$ for all $F'' \supset F'$. Define $F'_\alpha = F_\alpha \cup F'$. Then $[F'_\alpha] \supset [F_\alpha]$ and $G_k(\mu)_{[F'_\alpha]}$ has standard part which is ε -close to $G_k(\mu)$; moreover, every $[F''_\alpha] \supset [F'_\alpha]$ has the same property. It follows that the net of hyperfinite sums associated to $\text{cone}(F)$ have standard parts converging to $G_k(\mu)$. The infinitesimality statement follows. \square

The proof of the Proposition 9 reveals the function of cone ultrafilters: they are the ones that recognize classically convergent infinite series.

Define

$$\diamond\hat{G}_k^{\text{qt}} : \pm\mathbb{H} \times \mathbb{S}^1 \longrightarrow \diamond\hat{\Gamma}^{\text{qt}}, \quad (\mu, \theta) \mapsto \pi^{\text{qt}}([\theta])(\diamond\hat{G}_k(\mu)).$$

With the action of $\text{GL}_2(\mathbb{Z})$ on $\pm\mathbb{H} \times \mathbb{S}^1$ defined as in (21) we have

Proposition 10. *$\diamond\hat{G}_k^{\text{qt}}$ is a modular form of weight k :*

$$(A'(\mu))^k \cdot \diamond\hat{G}_k^{\text{qt}}(A(\mu, \theta)) = \diamond\hat{G}_k^{\text{qt}}(\mu, \theta).$$

Proof. Exactly the same proof as Proposition 8. \square

Let $\diamond\check{\mathbb{R}} \subset \diamond\check{\mathbb{C}}$ be the sheaf of real points, and denote by $\diamond\check{\Gamma}(\mathbb{R})$ the sections with values in $\diamond\check{\mathbb{R}}$. Let $\diamond\hat{\Gamma}^{\text{qt}}(\mathbb{R})$ denote the associated real points in $\diamond\hat{\Gamma}^{\text{qt}}$. For the value $\mu = i$, it is well-known that the classical Eisenstein series is real valued. For the same reasons we have the following important reality result for the classical Kronecker foliations (those corresponding to pairs (i, θ)):

Proposition 11. *For all k and $\theta \in \bar{\mathbb{R}}$, $\diamond\hat{G}_k^{\text{qt}}(i, \theta) \in \diamond\hat{\Gamma}^{\text{qt}}(\mathbb{R})$.*

Proof. As before, we work on the level of the defining net (25). We consider any subnet

$$\left\{ \sum_{[F_\alpha]} ({}^*n + {}^*n^\perp i)^{-2k} \right\}$$

where $[F_\alpha]$ range over the elements of some $X \in \mathfrak{c}(\theta)$. Taking the conjugate yields

$$\left\{ \sum_{[F_\alpha]} ({}^*n - {}^*n^\perp i)^{-2k} \right\} = \left\{ \sum_{[F_\alpha]} ({}^*n + {}^*n^\perp A(i))^{-2k} \right\}$$

where A is the element of $\text{PGL}_2(\mathbb{Z})$ defining $z \mapsto -z$. It follows then by the automorphy that $\diamond\hat{G}_k^{\text{qt}}(i, \theta)$ is equal to its own conjugate. \square

9. THE UNIVERSAL MODULAR INVARIANT

In this section we will define a universal modular invariant as a map of ultrasolenoids, in such a way that each of the classical and quantum invariants may be recovered from it as a subquotient (a restriction followed by quotients).

Define the **universal modular invariant**

$$\diamond \hat{j} : \pm\mathbb{H} \rightarrow \diamond \hat{\Gamma}$$

via the classical template [20]:

$$\diamond j(\mu) = 12^3 \cdot \frac{\diamond \hat{g}_2(\mu)^3}{\diamond \hat{g}_2(\mu)^3 - 27 \cdot \diamond \hat{g}_3(\mu)^2},$$

where the lower case (normalized) Eisenstein ultratransversals $\diamond \hat{g}_2, \diamond \hat{g}_3$ are defined in the usual way by scaling $\diamond \hat{G}_2, \diamond \hat{G}_3$ by 60 resp. 140.

The classical and quantum modular invariants are defined by composition with the projections π^{cl} resp. $\pi^{\text{qt}}(\theta)$ (see the end of §7), making the quantum invariant a function of $\pm\mathbb{H} \times \mathbb{S}^1$; since the automorphies of the numerator and denominator cancel, we obtain modular functions

$$\diamond j^{\text{cl}} := \pi^{\text{cl}} \circ \diamond \hat{j} : \text{Mod}^{\text{cl}} \longrightarrow \diamond \hat{\Gamma}^{\text{cl}}$$

and

$$\diamond j^{\text{qt}} : \text{Mod}^{\text{kf}} \longrightarrow \diamond \hat{\Gamma}^{\text{qt}}, \quad \diamond j^{\text{qt}}(\mu, \theta) = \pi^{\text{qt}}([\theta]) \circ \diamond j(\mu).$$

By Proposition 9, $\diamond \hat{g}_2^{\text{cl}} \simeq g_2$ and $\diamond \hat{g}_3^{\text{cl}} \simeq g_3$ so we have immediately:

Corollary 2. *Let $j(\mu)$ be the usual modular invariant of the elliptic curve $\mathbb{T}(\mu)$ viewed as a constant transversal in $\diamond \hat{\Gamma}^{\text{cl}}$. Then $\diamond \hat{j}^{\text{cl}}(\mu) \simeq j(\mu)$.*

Note that by Proposition 11, the image of $\text{Mod}_{[i]}^{\text{kf}}$ (the fiber over $[i]$) by $\diamond \hat{j}^{\text{qt}}$ belongs to the real locus:

Proposition 12. *For all $\theta \in \bar{\mathbb{R}}$, $\diamond \hat{j}^{\text{qt}}(i, \theta) \in \diamond \hat{\Gamma}^{\text{qt}}(\mathbb{R})$.*

The reality of the image of $\text{Mod}_{[i]}^{\text{kf}}$ given by Proposition 12 suggests that we may calculate $\diamond \hat{j}^{\text{qt}}(i, \theta)$ using hyperfinite partial sums over the group ${}^*\mathbb{Z}(\theta) \subset {}^*\mathbb{R}$ rather than over the group ${}^*\mathbb{A}(i, \theta) \subset {}^*\mathbb{C}$. This is reasonable, since by Proposition 6, every element of ${}^*\mathbb{A}(i, \theta)$ is of the form ${}^*n^\perp i + {}^*n$ for ${}^*n \in {}^*\mathbb{Z}(\theta)$.

We recall that the ultratransversal $\diamond \hat{j}^{\text{qt}}(i, \theta)$ is an equivalence class of the section $\diamond \hat{j}^{\text{qt}}(i, \theta)$ of the sheaf $\diamond \hat{\mathcal{C}}^{\text{qt}}$, where for each $u \in \text{Cone}(\mathcal{H})(\theta)$, the value of $\diamond \hat{j}^{\text{qt}}(i, \theta)$ is the u -class of the net of hyper-finite partial sums $\{j_{[F_\alpha]}\}$, for $[F_\alpha] \subset {}^*\mathbb{A}(i, \theta)$ hyper-finite. Let us write

$$j_{F_\alpha}(i, \theta) := \frac{12^3}{1 - J_{F_\alpha}(i, \theta)}, \quad J_{F_\alpha}(i, \theta) = \frac{49 \left(\sum_{mi+n \in F_\alpha} (mi+n)^{-6} \right)^2}{20 \left(\sum_{mi+n \in F_\alpha} (mi+n)^{-4} \right)^3}.$$

Since we are considering nets of hyperfinite partial sums which are increasing w.r.t. inclusion i.e. nets indexed by $X \in \mathfrak{c}(\theta)$, we may assume that the hyperfinite set $[F_\alpha]$ is symmetric w.r.t. multiplication by -1 , and write

$$J_{F_\alpha}(i, \theta) = \frac{49 \left(\sum_{mi+n \in F_\alpha, n>0} (mi+n)^{-6} \right)^2}{40 \left(\sum_{mi+n \in F_\alpha, n>0} (mi+n)^{-4} \right)^3}.$$

For such a hyperfinite set $[F_\alpha]$, let $[\bar{F}_\alpha] \subset {}^*\mathbb{Z}(\theta)$ be the set of ${}^*n \in {}^*\mathbb{Z}(\theta)$ for which ${}^*n^\perp i + {}^*n \in [F_\alpha]$. Consider the hyperfinite sum $j_{[\bar{F}_\alpha]}(\theta)$ defined as the * -class corresponding to the sequence

$$j_{F_\alpha}(\theta) := \frac{12^3}{1 - J_{\bar{F}_\alpha}(\theta)}, \quad J_{\bar{F}_\alpha}(\theta) = \frac{49 \left(\sum_{n \in \bar{F}_\alpha, n > 0} n^{-6} \right)^2}{40 \left(\sum_{n \in \bar{F}_\alpha, n > 0} n^{-4} \right)^3}.$$

Proposition 13. *There exists ${}^*u \in {}^*\mathbb{R}$, ${}^*u \simeq 1$, with*

$$j_{[F_\alpha]}(i, \theta) = {}^*u \cdot j_{[\bar{F}_\alpha]}(\theta).$$

In particular, $j_{[F_\alpha]}(i, \theta) \in {}^\mathbb{R}_{\text{fin}} \Leftrightarrow j_{[\bar{F}_\alpha]}(\theta) \in {}^*\mathbb{R}_{\text{fin}}$ and in this case, $j_{[F_\alpha]}(i, \theta) \simeq j_{[\bar{F}_\alpha]}(\theta)$.*

Proof. Let $[F_\alpha] \subset {}^*\Lambda(i, \theta)$; then we may write for each α

$$J_{F_\alpha}(i, \theta) = \frac{49 \left(\sum_{mi+n \in F_\alpha, n > 0} n^{-6} ((m/n)i + 1)^{-6} \right)^2}{40 \left(\sum_{mi+n \in F_\alpha, n > 0} n^{-4} ((m/n)i + 1)^{-4} \right)^3}.$$

Since $[F_\alpha] \subset {}^*\Lambda(i, \theta)$, for any $\varepsilon > 0$ we have $|\theta - m/n| = \varepsilon(n) < \varepsilon$ for all $mi + n \in F_\alpha$ and α sufficiently large. Multiplying numerator and denominator by $(\theta i + 1)^{12}$ and writing

$$(m/n)i + 1 = \theta i + 1 \pm \varepsilon(n)$$

gives the bounds

$$\left(\frac{\theta + 1}{\theta + 1 + \varepsilon} \right)^{12} J_{\bar{F}_\alpha}(\theta) < J_{F_\alpha}(i, \theta) < \left(\frac{\theta + 1}{\theta + 1 - \varepsilon} \right)^{12} J_{\bar{F}_\alpha}(\theta).$$

The result follows immediately. \square

Let ${}^\diamond \hat{\sigma} \in {}^\diamond \hat{\Gamma}^{\text{qt}}(\mathbb{R})$ be any section class. We say that ${}^\diamond \hat{\sigma}$ has standard part at (the $\text{GL}_2(\mathbb{Z})$ orbit of) u

$$\text{std}({}^\diamond \hat{\sigma}(u)) \in \mathbb{R}$$

if there exists a representative section ${}^\diamond \check{\sigma}$ such that for all $M \in \mathbb{R}_+$,

$$|{}^\diamond \check{\sigma}(u) - \text{std}({}^\diamond \hat{\sigma}(u))| < M.$$

Notice that if the standard part at u exists it is unique. If ${}^\diamond \hat{\sigma}$ does not have standard part at (the $\text{GL}_2(\mathbb{Z})$ orbit of) u we will write

$$\text{std}({}^\diamond \hat{\sigma}(u)) = \infty.$$

Thus each section class and each $\theta \in \mathbb{R}$ determines, in particular, a function on the $\text{GL}_2(\mathbb{Z})$ -orbit of $\text{Cone}(\mathcal{H})(\theta)$

$$\text{std}({}^\diamond \hat{\sigma})_\theta : \text{GL}_2(\mathbb{Z}) \setminus \left(\bigsqcup_{A \in \text{GL}_2(\mathbb{Z})} \text{Cone}(\mathcal{H})(A(\theta)) \right) \longrightarrow \bar{\mathbb{R}}.$$

This gives an induced multimap

$$\overline{\text{std}({}^\diamond \hat{\sigma})} : \mathbb{R} \dashrightarrow \bar{\mathbb{R}}, \quad \theta \mapsto \text{std}({}^\diamond \hat{\sigma})_\theta(\text{Cone}(\mathcal{H})(\theta)).$$

Theorem 7. *Let $\theta \in \mathbb{R} - \mathbb{Q}$. Then $\overline{\text{std}({}^\diamond j^{\text{qt}}(i, \theta))} \supset j^{\text{qt}}(\theta)$.*

Proof. Suppose that $j_0 \in j^{\text{qt}}(\theta)$ is the limit corresponding to the sequence $\{\varepsilon_\alpha\}$, whose class in ${}^*\mathbb{R}_\varepsilon$ is denoted ${}^*\varepsilon$. Consider a *shift function* $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ i.e. a function which is finite-to-1 and does not reverse the order: $\sigma(\alpha) \leq \sigma(\beta)$ if $\alpha \leq \beta$. For such a shift function, the sequence $\{\varepsilon_{\sigma(\alpha)}\}$ will produce the same limit j_0 . We denote the class of such a shifted sequence by $\sigma({}^*\varepsilon)$.² Note that if ${}^*\delta$ is any positive infinitesimal there exist shifts σ_0, σ_1 with $\sigma_0({}^*\varepsilon) < {}^*\delta < \sigma_1({}^*\varepsilon)$. Let

$${}^*\mathbb{Z}_{{}^*\varepsilon}^2(\theta) := \{({}^*m, {}^*n) \in {}^*\mathbb{Z}^2(\theta) \mid \pm {}^*n \in [B_{\varepsilon_\alpha}(\theta)]\}$$

where $[B_{\varepsilon_\alpha}(\theta)]$ is the ultraproduct of the $B_{\varepsilon_\alpha}(\theta) \subset \mathbb{N}$ defined in §1. We will produce a set of hyperfinites $X \subset \mathcal{H}$ compatible with the cone filter $\mathfrak{c}(\theta)$ (i.e. $X \cap \text{Cone}([F_\alpha]) \neq \emptyset$ for all $[F_\alpha] \subset {}^*\wedge(i, \theta)$) for which the net $\{j_{[F_\alpha]}\}_{[F_\alpha] \in X}$ converges to j_0 . Given $[F_\alpha] \subset {}^*\mathbb{Z}^2(\theta)$ we can find some shift map σ so that $[F_\alpha] \subset {}^*\mathbb{Z}_{\sigma({}^*\varepsilon)}^2(\theta)$. We may choose a hyperfinite $[\tilde{F}_\alpha] \subset {}^*\mathbb{Z}_{\sigma({}^*\varepsilon)}^2(\theta) \cap \text{Cone}([F_\alpha])$ such that $j_{[\tilde{F}_\alpha]} \simeq j_0$. Let X be the set of such $[\tilde{F}_\alpha]$. Then X is compatible with $\mathfrak{c}(\theta)$ so there exists a cone ultrafilter \mathfrak{u} containing X . This ultrafilter will produce a standard part j_0 . \square

In order to recover the invariant on the nose requires a slight paring down of $\text{Cone}(\theta)$. Consider the subset $J(\theta) \subset \mathcal{H}(\theta)$ of hyperfinites $[F_\alpha] \subset {}^*\mathbb{Z}^2(\theta)$ for which

$$j_{[F_\alpha]}(i, \theta) \simeq j_0 \in j^{\text{qt}}(\theta).$$

Clearly $J(\theta)$ intersects nontrivially every element of $\mathfrak{c}(\theta)$ (it intersects all the cones contained in $\mathfrak{c}(\theta)$), therefore we may form the filter $\mathfrak{c}(\theta)_0$ generated by $\mathfrak{c}(\theta)$ and $J(\theta)$. The set of ultrafilters \mathfrak{u} on \mathcal{H} extending $\mathfrak{c}(\theta)_0$ defines a closed subset

$$\text{Cone}(\theta)_0 \subset \text{Cone}(\theta).$$

Note that $\text{Cone}(\theta)_0$ is taken to $\text{Cone}(A(\theta))_0$ by any $A \in \text{GL}_2(\mathbb{Z})$, since $A(J(\theta)) = J(A(\theta))$: the latter follows from the fact that the standard j^{qt} is $\text{GL}_2(\mathbb{Z})$ -invariant. If we denote by ${}^\diamond j_0^{\text{qt}}(i, \theta)$ the restriction of ${}^\diamond j^{\text{qt}}(i, \theta)$ to $\text{Cone}(\theta)_0$ then we have

Theorem 8. *Let $\theta \in \mathbb{R} - \mathbb{Q}$. Then $\overline{\text{std}}({}^\diamond j_0^{\text{qt}}(i, \theta)) = j^{\text{qt}}(\theta)$.*

Proof. By the previous Theorem we only need to show that every point in $\overline{\text{std}}({}^\diamond j_0^{\text{qt}}(i, \theta))$ belongs to $j^{\text{qt}}(\theta)$. If j_0 is a standard part of ${}^\diamond j_0^{\text{qt}}(i, \theta)$ converging with respect to the restriction to $X \in \mathfrak{u}$, then we may produce an element $[F_\alpha] \in J(\theta) \cap X$ with $j_{[F_\alpha]}(i, \theta) \simeq j_0$ and by definition of $J(\theta)$, this shows that $j_0 \in j^{\text{qt}}(\theta)$. \square



We finish with an equivalent formulation of ${}^\diamond j$ that evokes the framework of the classical modular invariant. Let

$$\pm {}^\diamond \mathbb{H} \subset {}^\diamond \check{\mathbb{C}}$$

be the subsheaf of ${}^\diamond \check{\mathbb{C}}$ having stalk $\pm {}^\diamond \mathbb{H}_u \subset {}^\diamond \mathbb{C}_u$ for each $u \in \text{Ult}(\mathcal{H})$. On $\pm {}^\diamond \mathbb{H}$ we define a diagonal action of $\text{GL}_2(\mathbb{Z})$, given, at the level of nets, by

$$\{z_{[F_\alpha]}\} \mapsto \{w_{[F_\alpha]}\} := \{A(z_{[A^T F_\alpha]})\}.$$

The action of A is therefore a shifted linear action. We denote it by

$${}^\diamond z_u \mapsto A \otimes {}^\diamond z_u \in \mathbb{C}_{A^{-T}u}$$

²Note that the map σ does not induce a function on ${}^*\mathbb{R}$.

to distinguish it from the earlier defined $A \circledast z_u$, the shift induced by A . The quotient by this action is an ultrasolenoid denoted

$$\circledast \widehat{\text{Mod}},$$

whose «leaves» are the images of the stalks of $\pm \circledast \check{\mathbb{H}}$.

The action which defines $\circledast \widehat{\text{Mod}}$ is the analogue of the diagonal action of $\text{GL}_2(\mathbb{Z})$ on $\pm \mathbb{H} \times \mathbb{S}^1$ used to define the signed Anosov foliation Mod^{kf} . We may think of each point of $\circledast \widehat{\text{Mod}}$ as parametrizing an isomorphism class of abstract nonstandard Kronecker foliation in the nonstandard elliptic curve

$$\circledast \mathbb{C}_u / \circledast \Lambda(\circledast z_u),$$

where $\circledast \Lambda(\circledast z_u)$ is the $\circledast \mathbb{Z}_u$ -module generated by $1, \circledast z_u$ and where moreover each $u \in \text{Ult}(\mathcal{H})$ is to be thought of as supplying the data of an «ultraslope».

Denote by $\circledast \text{Mod}$ the image in $\circledast \widehat{\text{Mod}}$ of the constant sheaf $\pm \mathbb{H}$ over $\text{Ult}(\mathcal{H})$ with stalk $\pm \mathbb{H} \subset \pm \circledast \mathbb{H}_u$ for all $u \in \text{Ult}(\mathcal{H})$. The image of $\pm \mathbb{H}^{\text{cl}}$ (= the restriction of $\pm \mathbb{H}$ to $\text{Cone}(\mathcal{H})$) in $\circledast \text{Mod}$ is denoted $\circledast \text{Mod}^{\text{cl}}$; likewise, the image of the restriction to the union of the θ -quantum cone ultrafilters is denoted $\circledast \text{Mod}^{\text{qt}}$.

Given a hyperfinite set $[F_\alpha] \in \mathcal{H}$ and ${}^* z_{[F_\alpha]} = {}^* \{z_{F_\alpha}\} \in \pm {}^* \mathbb{H} \subset {}^* \mathbb{C}$, let

$$G_k({}^* z_{[F_\alpha]}) := {}^* \left\{ \sum_{(m,n) \in F_\alpha} (m z_{F_\alpha} + n)^{-2k} \right\}.$$

Following the usual procedure we may then define $j({}^* z_{[F_\alpha]})$. Extending to a map of nets indexed by \mathcal{H} leads to a function of sheaves

$$\circledast j : \pm \circledast \check{\mathbb{H}} \longrightarrow \circledast \check{\mathbb{C}}.$$

In the Theorem which follows we will need to specify topologies on the various sheaves and ultrasolenoids defined above. This can be done by putting a topology on $\circledast \check{\mathbb{C}}$ as follows. First, we topologize each fiber $\circledast \mathbb{C}_u$ using the $(\circledast \mathbb{R}_u)_+$ -valued absolute value $\circledast |\cdot|_u$. Note that there is a canonical inclusion of the set of \mathcal{H} -nets

$${}^* \mathbb{C}^{\mathcal{H}} \hookrightarrow \circledast \check{\mathbb{C}}$$

given by

$$\{{}^* z_{[F_\alpha]}\} \mapsto (u \mapsto \circledast \{{}^* z_{[F_\alpha]}\}_u).$$

We denote this section simply $\circledast z$, and its value at u by $\circledast z_u$. Now given $\mathcal{O} \subset \text{Ult}(\mathcal{H})$, $\{{}^* z_{[F_\alpha]}\} \subset {}^* \mathbb{C}$ and $\{{}^* r_{[F_\alpha]}\} \subset {}^* \mathbb{R}_+$ a pair of \mathcal{H} -nets, let

$$\check{\mathcal{O}}(\circledast z; \circledast r) = \{\circledast w_u \mid u \in \mathcal{O}, |\circledast z_u - \circledast w_u|_u < \circledast r_u\}.$$

The sets $\check{\mathcal{O}}(\circledast z; \circledast r)$ form the base for a topology on $\circledast \check{\mathbb{C}}$, called the **ultrasheaf topology**. Observe that the subspace topology on $\mathbb{C} \times \text{Ult}(\mathcal{H}) \subset \circledast \check{\mathbb{C}}$ coincides with the product topology.

Note that any section $\circledast w$ defined by a net $\{{}^* w_{[F_\alpha]}\}$ is continuous with respect to the ultrasheaf topology: indeed, if $\check{\mathcal{O}}(\circledast z; \circledast r)$ contains the point $\circledast w_u$, then there exists a subset $X \in u$ such that the subnet $\{{}^* w_{[F_\alpha]}\}_X$ is contained in the subnet of balls $\{B_{{}^* r_{[F_\alpha]}}({}^* z_{[F_\alpha]})\}_X$, where $B_{{}^* r_{[F_\alpha]}}({}^* z_{[F_\alpha]})$ is the ball of radius ${}^* r_{[F_\alpha]}$ about ${}^* z_{[F_\alpha]}$. But this implies that this is true for any ultrafilter u' containing X . In other words, the pre-image of the open $\check{\mathcal{O}}(\circledast z; \circledast r)$ is the union of the Stone opens $\mathcal{O}_X = \{u \ni X\}$, where X is as above.

More generally, any map $\circledast \check{\mathbb{C}} \longrightarrow \circledast \check{\mathbb{C}}$ which takes stalks to stalks is continuous if it is continuous along the base and if each map $\circledast \mathbb{C}_u \rightarrow \circledast \mathbb{C}_{u'}$ is continuous in the

norm topologies In particular, both the shift and the diagonal actions of $\mathrm{GL}_2(\mathbb{Z})$ on $\pm^\circ\mathbb{H} \subset {}^\circ\hat{\mathbb{C}}$ act by homeomorphisms, properly discontinuously, so that the induced quotient topologies on ${}^\circ\hat{\mathbb{C}}$ – as well as on ${}^\circ\widehat{\mathrm{Mod}}$ and each of its subsolenoids – are Hausdorff. Note that on ${}^\circ\mathrm{Mod} \subset {}^\circ\widehat{\mathrm{Mod}}$, the subspace topology coincides with the topology induced by the product topology $\mathbb{H} \times \mathrm{Ult}(\mathcal{H})$.

Theorem 9. *The map of nets $\{z_{[F_\alpha]}\}_{[F_\alpha] \in \mathcal{H}} \mapsto \{j^*(z_{[F_\alpha]})\}_{[F_\alpha] \in \mathcal{H}}$ induces a continuous leaf-preserving map of ultrasolenoids*

$${}^\circ j : {}^\circ\widehat{\mathrm{Mod}} \longrightarrow {}^\circ\hat{\mathbb{C}}.$$

The restriction of ${}^\circ j$ to ${}^\circ\mathrm{Mod}$ is induced by the universal modular invariant ${}^\circ\hat{j} : \pm\mathbb{H} \rightarrow {}^\circ\hat{\Gamma}$ by the formula

$${}^\circ j(z, u) = {}^\circ\hat{j}(z)(u).$$

In particular, if we denote by ${}^\circ j^{\mathrm{cl}}, {}^\circ j^{\mathrm{qt}}$ the restriction of ${}^\circ j$ to ${}^\circ\mathrm{Mod}^{\mathrm{cl}}$ resp. ${}^\circ\mathrm{Mod}^{\mathrm{qt}}$ then

$${}^\circ j^{\mathrm{cl}}(z, u) = {}^\circ\hat{j}^{\mathrm{cl}}(z)(u), \quad {}^\circ j^{\mathrm{qt}}(z, u) = {}^\circ\hat{j}^{\mathrm{qt}}(z)(u).$$

Proof. Given $A \in \mathrm{GL}_2(\mathbb{Z})$, define $A' \otimes {}^\circ z_u \in {}^\circ\mathbb{C}_{A^{-T}u}$ via the action on representative nets:

$$\{A'({}^*z)_{[A^T F_\alpha]}\}.$$

Then the map of sheaves ${}^\circ G_k : \pm^\circ\mathbb{H} \rightarrow {}^\circ\hat{\mathbb{C}}$ satisfies

$$(A' \otimes {}^\circ z_u)^k \cdot {}^\circ G_k(A \otimes {}^\circ z_u) = A \circ {}^\circ G_k({}^\circ z_u).$$

From this, it follows immediately that ${}^\circ j$ induces a function ${}^\circ j : {}^\circ\widehat{\mathrm{Mod}} \rightarrow {}^\circ\hat{\mathbb{C}}$, and that the functions ${}^\circ j, {}^\circ j^{\mathrm{cl}}, {}^\circ j^{\mathrm{qt}}$ are related to ${}^\circ\hat{j}, {}^\circ\hat{j}^{\mathrm{cl}}, {}^\circ\hat{j}^{\mathrm{qt}}$ by the formulas given in the statement of the Theorem. If we fix a net $\{z_{[F_\alpha]}\}$, the association $u \mapsto {}^\circ z_u$ gives a continuous section of $\pm^\circ\mathbb{H}$ and therefore $u \mapsto {}^\circ j({}^\circ z_u)$ is continuous as well. If we fix $u \in \mathrm{Ult}(\mathcal{H})$ then the map $\pm^\circ\mathbb{H}_u \rightarrow {}^\circ\mathbb{C}_u, {}^\circ z_u \mapsto {}^\circ j({}^\circ z_u)$ gives a continuous map of the stalk ${}^\circ\mathbb{C}_u$: by transference from the continuity of the classical modular invariant. Thus ${}^\circ j$ is continuous in the ultrasheaf topology. \square

Note the dual nature of the classical and quantum invariant: the values of the classical invariant are recovered *along the leaves* of a restriction, whereas the quantum invariant is recovered *along the transversal* of another.

10. APPENDIX: EXPERIMENTAL VALUES OF J^{qt}

In this section we present some preliminary experimental evidence which suggests that for $\theta \in \mathbb{R}$ a quadratic irrationality, $J^{\mathrm{qt}}(\theta)$ is a finite set.

Let D be a fundamental discriminant. For u the fundamental unit in $\mathbb{Q}(\sqrt{D})$ the function

$$x \mapsto J_{u^{-x}}^{\mathrm{qt}}(u)$$

is mapped using the plth function of PARI/GP. Below are the results for the first five fundamental discriminants $D = 5, 8, 12, 13$ and 17 .

In the case of discriminant 5, the experimental lower bound 0.8188... matches the value obtained by calculating the formula (12) and gives $J_{\mathrm{best}}^{\mathrm{qt}}(\varphi) \approx 9538.2496...$ On the other hand, if we consider the modification of (12) obtained by replacing $G_M(\varphi)$ by $G'_M(\varphi) = G_M(\varphi) + 1$, a value very close to 0.8501... = the experimental supremum of $J^{\mathrm{qt}}(\varphi)$ is returned.

More generally, close examination of the graphs below suggests that the number of returned values of $J^{\mathrm{qt}}(u)$ is approximately D . Based on these (very preliminary)

computations it doesn't seem entirely unreasonable to contemplate the following rough

Conjecture. *Let $\theta \in \mathbb{R} - \mathbb{Q}$ be a quadratic irrationality belonging to $\mathbb{Q}(\sqrt{D})$. Then $j^{qt}(\theta)$ is a finite bounded set. If $\theta = u$ is a fundamental unit then*

$$\text{card}(j^{qt}(u)) = O(D).$$




FIGURE 1. $D=5$.







FIGURE 2. $D=8$.

FIGURE 3. $D=12$.FIGURE 4. $D=13$.FIGURE 5. $D=17$.

REFERENCES

- [1] Baxter, R.J., *Exactly Solved Models in Statistical Mechanics*. Reprint of the 3rd edition of the work originally published by Academic Press. Dover Books on Physics, Dover Publications, Mineola, NY, 2008.
- [2] Cassels, J.W.S. *An introduction to Diophantine approximation*. Cambridge Tracts in Mathematics and Mathematical Physics, **45**. Cambridge University Press, New York, 1957.
- [3] Chang, C.C. & Keisler, H.J., *Model Theory*. 3rd Ed. Studies in Logic and the Foundations of Mathematics **73**, North Holland, Amsterdam, 1990.
- [4] Connes A. & Marcolli, M. *Noncommutative Geometry, Quantum Fields and Motives*, AMS Colloquium Publications **55**, AMS, Providence, RI, 2008.
- [5] Connes A., Marcolli, M. & Ramachandran, N., KMS states and complex multiplication I, *Selecta Math.* (New Ser.) **11** (2005), no. 3-4, 325-347.
- [6] Connes A., Marcolli, M. & Ramachandran, N., KMS states and complex multiplication II, in "Operator Algebras: The Abel Symposium 2004", pp. 15-59, Abel Symp. **1**, Springer-Verlag, New York, 2006.
- [7] Dasgupta, S., Darmon, H. & Pollack, R., Hilbert modular forms and the Gross-Stark conjecture. *Ann. Math.* **174**, no. 1 (2011), 439-484.
- [8] Eklov, P.C., Ultraproducts for algebraists. in *Handbook of Mathematical Logic* (J. Barwise, ed.), Studies in Logic and the Foundations of Mathematics, pp. 105-137, North Holland, Amsterdam, 1989.
- [9] Fesenko, I.B., Model theory guidance in number theory? In "Model Theory with Applications to Algebra and Analysis", LMS Lecture Note Series **349**, pp. 327-334 eds. Z. Chatzidakis, D. Macpherson, A. Pillay, A. Wilkie, Cambr. Univ. Press, Cambridge U.K., 2008.
- [10] Fesenko, I.B., Several nonstandard remarks, In "AMS/IP Advances in the Mathematical Sciences", AMS Transl. Series 2, **217** (2006), 37-50.
- [11] Gendron, T.M., The algebraic theory of the fundamental germ. *Bull. Braz. Math. Soc. (N.S.)* **37** (2006), no. 1, 49-87.
- [12] Gendron, T.M., The geometric theory of the fundamental germ. *Nagoya Math. J.* **190** (2008), 1-34.
- [13] Gendron, T.M., Real algebraic number theory I: Diophantine approximation groups, Kronecker foliations and independence. arXiv:1201.2708v1 (2012).
- [14] Gendron, T.M., Real algebraic number theory II: Ideological arithmetic. arXiv:1208.4334v2 (2012).
- [15] Gendron, T.M., Real algebraic number theory III: Transcendental Galois theory. *in preparation*.
- [16] Goldblatt, R., *Lectures on the Hyperreals: An Introduction to Nonstandard Analysis*. Graduate Texts in Mathematics **188**, Springer-Verlag, 1998.
- [17] Hardy, G.H. & Wright, E.M., *An Introduction to the Theory of Numbers*. 5th edition. Oxford Science Publications, Oxford University Press, Oxford, UK, 1980.
- [18] Hecke, E., Höhere Modulfunktionen und ihre Anwendung auf die Zahlentheorie. *Math. Ann.* **71** (1912), 1-37.
- [19] Hodges, W., *Model Theory*. Encyclopedia of Mathematics and its Applications **42**. Cambridge University Press, Cambridge, 1993.
- [20] Husemöller, D., *Elliptic Curves*, 2nd Ed., Graduate Texts in Mathematics **111**, Springer-Verlag, New York, 2004.
- [21] Jech, T., *Set Theory*. Third Millennium Edition. Springer-Verlag, New York, 2006.
- [22] Johnstone, T., *Stone Spaces*. Cambridge Studies in Advanced Mathematics, Vol. 3, Cambridge University Press, New York, 1983.
- [23] Kaneko, M., Observations on the 'values' of the elliptic modular function $j(\tau)$ at real quadratics. *Kyushu J. Math.* **63** (2009), 353-364.
- [24] MacIntyre, A., Nonstandard analysis and cohomology. Nonstandard methods and applications in mathematics, 174-191, Lect. Notes Log., 25, Assoc. Symbol. Logic, La Jolla, CA, 2006.
- [25] Manin, Yu. I., Real multiplication and noncommutative geometry. in "The Legacy of Niels Henrik Abel", pp. 685-727, Springer-Verlag, New York, 2004.
- [26] Manin, Yu. I., An update on real multiplication and noncommutative geometry. http://www.hcm.uni-bonn.de/fileadmin/user_upload/Ellipticintegrablesystems/Manin.pdf.
- [27] Marcolli, M., *Arithmetic Noncommutative Geometry*, University Lecture Series **36**, AMS, Providence, RI, 2005.
- [28] Marker, D., *Model Theory: An Introduction*, Graduate Texts in Mathematics **217**, Springer-Verlag, New York, 2002.
- [29] Niven, I., Zuckerman, H.S. & Montgomery, H.L., *An Introduction to the Theory of Numbers*, 5th edition, John Wiley & Sons, Inc., New York, 1991.

- [30] Ostrowski, V. A., Bemerkungen zur theorie der diophantischen approximationen, *Abh. Math. Semin. Hamburg Univ.* **1** (1922), 77–98.
- [31] Poizat, B., *A Course in Model Theory*. (trans. by Moses Klein). Universitext, Springer-Verlag, New-York, 2000.
- [32] Robinson, A., *Non-standard Analysis*. Princeton Landmarks in Mathematics, Princeton U. Press, Princeton NJ, 1996.
- [33] Schmidt, W.M., *Diophantine Approximation*. Lecture Notes in Mathematics **785**, Springer-Verlag, New York, 1980.
- [34] Serre, J.-P., Complex multiplication. in ‘Algebraic Number Theory: Proceedings of an Instructional Conference Organized by the London Mathematical Society’ (ed. by J. W. S. Cassels and Fröhlich), pp. 190–197.
- [35] Shimura, G., Class fields over real quadratic fields and Hecke operators. *Ann. of Math.* (2) **95** (1972), 130–190.
- [36] Silverman, J.H. *Advanced Topics in the Arithmetic of Elliptic Curves*. Graduate Texts in Mathematics **151**, Springer-Verlag, New York, 1986.
- [37] Stark H.M., Hilbert’s twelfth problem and L-series. *Bull. Am. Math. Soc.* **83**, No.5 (1977), 1072–1074.
- [38] Taylor, L., A Nonstandard Approach to Real Multiplication. arXiv/math/0612184.
- [39] Taylor, L., Higher Derivatives of L-series associated to Real Quadratic Fields. arXiv/math/0612186.
- [40] Taylor, L., Line Bundles over Quantum Tori. arXiv/math.NT/0612189.
- [41] Vorobiev, Nicolai N., *Fibonacci Numbers*, Birkhäuser Verlag, Basel, 2002.
- [42] Zeckendorf, E., Représentation des nombres naturels par une somme de nombres de Fibonacci ou de nombres de Lucas. *Bull. Soc. R. Sci. Liège* **41** (1972), 179–182.

CENTRO DE ESTUDIOS EN FÍSICA Y MATEMÁTICAS BÁSICAS Y APLICADAS, UNIVERSIDAD AUTÓNOMA DE CHIAPAS, 4A. ORIENTE NORTE NO. 1428, COLONIA BARRIO LA PIMIENTA, TUXTLA GUTIÉRREZ, CHIAPAS, MÉXICO

E-mail address: ccastanobernard@gmail.com

INSTITUTO DE MATEMÁTICAS – UNIDAD CUERNAVACA, UNIVERSIDAD NACIONAL AUTONOMA DE MÉXICO, AV. UNIVERSIDAD S/N, C.P. 62210 CUERNAVACA, MORELOS, MÉXICO

E-mail address: tim@matcuer.unam.mx